

Volume 6, No. 6, November 2016 * Bimonthly

ISSN: 2251-9939

Journal of Life Science and Biomedicine



Available online at www.jlsb.science-line.com

*Published by
Scienceline Publications*



Editorial Team

Editor-in-Chief: Parham Jabbarzadeh Kaboli

PhD of Molecular Biology and Cancer researcher; Faculty of Medicine and Health Sciences, University Putra, Malaysia ([Website](#); [Emails: researchgroups@drugdiscovery.ir](#))

Managing Editor: Yusuf Kaya

PhD, Professor of Biology, Atatürk University, Erzurum, ([Website](#), [Email: ykaya@atauni.edu.tr](#))

Executive Editor: Zohreh Yousefi

PhD candidate, Biosystematics, Atatürk University, Erzurum, Turkey ([Emails: zohreh.yousefi12@ogr.atauni.edu.tr](#))

Language Editor: Samuel Stephen Oldershaw

Master of TESOL, The Humberston School & The Grimsby Institute, Nuns Corner, Grimsby, North East Lincolnshire, United Kingdom ([Email: s.s.oldershaw@hotmail.com](#))

Associate Editors

Aleksandra K. Nowicka

PhD, Pediatrics and Cancer researcher; MD Anderson Cancer Center, Houston, Texas, USA ([Email: aknowicka@mdanderson.org](#))

Paola Roncada

PhD, Pharmacokinetics, Residues of mycotoxins in food and in foodproducing species, University of Bologna, Italy ([Email: paola.roncada@unibo.it](#))

Tohid Vahdatpour

PhD, Assistant Prof., Physiology, Islamic Azad University, Iran ([Website](#); [Scopus](#); [Emails: vahdatpour@iaushab.ac.ir](#))

Veghar Hejazi

MD, Tabriz University of Medical Sciences, Tabriz, Iran ([Email: vegharhejazi@gmail.com](#))

Nefise Kandemir

MD, PhD, Department of Medical Genetics, Erciyes University, Kayseri, Turkey

Reviewers

Abolghasem Yousefi

PhD, Assistant Professor of Anesthesiology, Tehran University of Medical Sciences, Tehran, Iran ([Website](#); [Email: ayousefi@gmail.com](#))

Aleksandra K. Nowicka

PhD, Pediatrics and Cancer researcher; MD Anderson Cancer Center, Houston, Texas, USA ([Email: aknowicka@mdanderson.org](#))

Amany Abdin

PhD, Pharmacology; MSc, Medical Biochemistry; Tanta University, Egypt ([Emails: amanyabdin@med.tanta.edu.eg, amanyahr@hotmail.com](#))

Babak yousefi

Physician, General Surgery Resident at Hamedan University of Medical Science, Hamedan, Iran

Fazal Shirazi

PhD, Infectious Disease researcher at MD Anderson Cancer Center, Houston, Texas, USA

Fikret Çelebi

Professor of Veterinary Physiology; Atatürk University, Turkey ([Website](#); [Email: fncelebi@atauni.edu.tr](#))

Ghada Khalil Al Tajir

PhD, Pharmacology, Faculty of Medicine, UAE University, Al Ain, UAE

M.R. Ghavamnasiri

PhD, Professor of Oncology at Omid Cancer Hospital, MUMS; Cancer Research Center, Mashhad University of Medical Sciences, Iran

Kaviarasan Thanamegm

PhD of Marine Bioactive compounds, Department of Ecology and Environmental Sciences, Pondicherry University, India ([Email: marinekavi@gmail.com](mailto:marinekavi@gmail.com))

Jahan Ara Khanam

PhD, Anti-cancer Drug Designer and Professor of UR; Department of Biochemistry and Molecular Biology, University of Rajshahi, Bangladesh

Mozafar Bagherzadeh Homaee

PhD, Plant Physiology, University of Isfahan, Isfahan, Iran

Osman Erganiş

Professor, PhD, Veterinary Microbiology, Selcuk University, Konya, Turkey

Paola Roncada

PhD, Pharmacokinetics, Residues of mycotoxins in food and in foodproducing species, University of Bologna, Italy ([Email: paola.roncada@unibo.it](mailto:paola.roncada@unibo.it))

Perumal Karthick

Professor, PhD, Marine Biology, Pondicherry University, Brookshabad Campus, Port Blair, Andamans. 744112, India ([Email: karthickmicrobes@gmail.com](mailto:karthickmicrobes@gmail.com))

Reza Khodarahmi

PhD, Biochemistry at KU; Pharmacy School, Kermanshah University, Kermanshah, Iran

Saeid Chekani Azar

PhD, Veterinary Physiology, Atatürk University, Erzurum, Turkey ([Google Scholar](https://scholar.google.com/citations?user=saeid.azar); [Emails: saeid.azar@atauni.edu.tr](mailto:saeid.azar@atauni.edu.tr); schekani@gmail.com)

Siamk Sandoughchian

PhD Student, Immunology, Juntendo University, Japan

Siva Sankar. R.

PhD, Marine Biology, Dept. of Ecology & Environmental Sciences, Pondicherry University, Puducherry - 605014, India ([Email: sivauniverse@gmail.com](mailto:sivauniverse@gmail.com))

Tohid Vahdatpour

PhD, Assistant Prof., Physiology, Islamic Azad University, Iran ([Website](#); [Scopus](#); [Google Scholar](#); [Emails: vahdatpour@iaushab.ac.ir](mailto:vahdatpour@iaushab.ac.ir))

Veghar Hejazi

MD, Tabriz University of Medical Sciences, Tabriz, Iran ([Email: vegharhejazi@gmail.com](mailto:vegharhejazi@gmail.com))

Yusuf Kaya

PhD, Professor of Plant Biology, Atatürk University, Erzurum, Turkey ([Email: ykaya@atauni.edu.tr](mailto:ykaya@atauni.edu.tr))

Volume 6 (6); November 25, 2016

Research Paper

Handling, Processing and Utilization of Milk and Its Products in Gondar Town, Ethiopia.


Addisu Sh, Muhammed A and Haile N.
J. Life Sci. Biomed., 6 (6): 120-126, 2016;
 pii:S225199391600020-6

Abstract

The study was conducted from November to May, 2016 with the objective to assess the handling, processing and marketing of milk and its product in Gondar town. Data were collected with the three representative kebeles (kebele 18, 19 and 20) purposively. The study was carried out through informal and formal surveys. From those kebeles a total of 45 respondents were taken randomly. From the respondents majority of them were male (86.7%). 42.2% of the respondents were keep dairy cow under the age of 47-62. In the study area, the overall average family size was 5.7%. The equipments used for milking were gourd (28.9%), plastic jar (64.4%) and aluminium jar (6.67%). Equipments used for milk processing were gourd (48.89%), cream separator (15.5%), clay pot (11.11%) and others (24.4%). More than half of the respondents had not used refrigerator for handling of milk and its products. All milkers were washed their hands before milking however, only 62% of the respondents were washed the cow udder before milking. 60% of the respondents were processed the milk in traditional way. Preferred milk processed products were skim milk (48.89%), butter (40%), yogurt (6.67%) and the remaining were cheese (4.44%). Half of the respondents were used yogurt for household consumption however, 77.77% of the respondents were selling milk in the market. Generally milk production in Gondar town is contributing a role for the household livelihood improvement however, handling, processing and utilization of milk and its products should need further improvement.

Keywords: Gondar, Handling, Processing, Utilization of Milk

[Full text-[PDF](#)] [[XML](#)]



Research Paper

Phenotypic Characterization of Indigenous Cattle Populations in West Gojjam Administrative Zones, Amhara National Regional State, Ethiopia.

Tenagne A, Mekuriaw G and D Kumar.
J. Life Sci. Biomed., 6 (6): 127-138, 2016;
 pii:S225199391600021-6

Abstract

The study was carried out in Semen Achefer, Sekela and Jabitenan districts of Western Gojjam zone of Amhara National Regional State. The objective of the study was to carry out phenotypic characterization of local cattle population in the study under farmers' management condition in the study area. A total of 600 cattle were sampled randomly for characterization of phenotypic traits. Data were gathered through field observations and linear body measurements of sample populations. The Sampled indigenous cattle were identified by sex and district (agro ecology). The most dominant coat colour patterns in the sampled populations were plain, patchy and spotted with the most frequently observed coat colour type being light red, black and dark red. Sex of animals had $P < 0.05$, on all of the body measurements. Agro ecology also showed $P < 0.05$, for most of the body measurements, except tail length, horn length, height at wither and rump height. Moderate and $P < 0.05$, positive correlation was found among the body measurements. The prediction of body weight could be based on regression equation $y = -481.55 + 4.89x$ for male sample population and $y = -405.22 + 4.64x$ for female sample cattle population where y and x are body weight and chest girth, respectively. Most of the body measurements of cattle were affected by sex and agro- ecology. Phenotypic result of cattle populations in the study areas was varied from former finding and therefore; to put specific characteristics' of the breed, further molecular characterization is needed.

Keywords: Body Weight, Cattle, Characterization, Indigenous, Linear Body Measurement

[Full text-[PDF](#)] [[XML](#)]



Journal of Life Science and Biomedicine



ISSN: 2251-9939

Frequency: Bimonthly

Current Issue: 2016, Vol: 6, Issue 6 (November)

Publisher: [SCIENCELINE](#)

The Journal of Life Science and Biomedicine is aimed to improve the quality and standard of life with emphasis on the related branches of science such as biology, physiology, biochemistry, zoology, anatomy, pathology and their applications and innovations in medicine and healthcare... [view full aims and scope](#)

<http://jlsb.science-line.com>

» JLSB indexed/covered by [NLM Catalog](#), [RiCeST \(ISC\)](#), [Ulrich's™](#), [SHERPA/RoMEO](#), [Genamics](#), [Google Scholar \(h-index= 10\)](#), [Index Copernicus](#), [ICV2015: 66.26...](#) [details](#)

» Open access full-text articles is available beginning with Volume 1, Issue 1.

» Full texts and XML articles are available in ISC-RiCeST.

» This journal is in compliance with [Budapest Open Access Initiative](#) and [International Committee of Medical Journal Editors' Recommendations](#).



» High visibility of articles over the internet.

» Publisher Item Identifier [...details](#)

» This journal encourage the academic institutions in low-income countries to publish high quality scientific results, free of charges... [view Review/Decisions/Processing/Policy](#)



[ABOUT US](#)

| [CONTACT US](#)

| [PRIVACY POLICY](#)

Editorial Offices:

Atatürk University, Erzurum 25100, Turkey

University of Manitoba, Winnipeg, Manitoba R3T 2N2, Canada

University of Maragheh, East Azerbaijan, Maragheh 55136, Iran

Homepage: www.science-line.com

Phone: +98 914 420 7713 (Iran); +90 538 770 8824 (Turkey); +1 204 8982464 (Canada)

Emails:

administrator@science-line.com

saeid.azar@atauni.edu.tr

Handling, Processing and Utilization of Milk and Its Products in Gondar Town, Ethiopia

Shewangzaw Addisu^{1✉}, Ahmed Muhammed and Nunu Haile

University of Gondar, Faculty of Veterinary Medicine P.O.Box, 196, Gondar, Ethiopia

✉Corresponding author's Email: shewangzaw21@yahoo.com

Received 28 Sep. 2016 Accepted 05 Nov. 2016

ABSTRACT: The study was conducted from November to May, 2016 with the objective to assess the handling, processing and marketing of milk and its product in Gondar town. Data were collected with the three representative kebeles (kebele 18, 19 and 20) purposively. The study was carried out through informal and formal surveys. From those kebles a total of 45 respondents were taken randomly. From the respondents majority of them were male (86.7%). 42.2% of the respondents were keep dairy cow under the age of 47-62. In the study area, the overall average family size was 5.7%. The equipments used for milking were gourd (28.9%), plastic jar (64.4%) and aluminum jar (6.67%). Equipments used for milk processing were gourd (48.89%), cream separator (15.5%), clay pot (11.11%) and others (24.4%). More than half of the respondents had not used refrigerator for handling of milk and its products. All milkers were washed their hands before milking however, only 62% of the respondents were washed the cow udder before milking. 60% of the respondents were processed the milk in traditional way. Preferred milk processed products were skim milk (48.89%), butter (40%), yogurt (6.67%) and the remaining were cheese (4.44%). Half of the respondents were used yogurt for household consumption however, 77.77% of the respondents were selling milk in the market. Generally milk production in Gondar town is contributing a role for the household livelihood improvement however, handling, processing and utilization of milk and its products should need further improvement.

Author Keywords: Gondar, Handling, Processing, Utilization of Milk

INTRODUCTION

Demand for milk and dairy products has increased in the tropical areas where peoples growing. However, despite milk's contribution to gross domestic products and its value as a food, sub-Saharan Africa in general and Ethiopia in particular has failed to attain self-sufficiency in dairy production [1]. Dairy processing plants are few in numbers, much of the milk produced by rural small holders is processed on farm using traditional technologies and milk processing is based on sour milk [2]. This is due to high ambient temperature, small daily quantities of milk produced, consumer preference, the better shelf life of fermented milk as well as the type and capacity of the locally available processing materials and methods used. Ethiopians, like other countries, have been using milk as part of their diet for centuries. Milk is used for rearing calves and children and the surplus is soured for Ergo (Ethiopian naturally fermented milk) and/or butter and Ayib (Ethiopian cottage cheese) making. Arera (defatted sour milk) is used for human consumption or for Ayib-making.

In Ethiopia, there is no standard hygienic condition followed by producers during milk production. The hygienic conditions are different according to their production system. In most of the cases under smallholder condition, the common hygienic measures taken during milk production especially during milking are limited to letting the calf to suckle for few minutes and/or washing the udder before milking. The quality of the water used for cleaning purpose (washing the udder, milk equipment, hands), however, is not secured [3]. Milk production under Ethiopian smallholder condition can therefore generally be characterized as free from poisoning. Dairy products are manufactured and consumed in many parts of the country. Fresh whole milk, whole sour milk (Ergo), butter, Arera (defatted sour milk) and Ayib (a traditional cottage cheese) are the major dairy products produced and consumed in many parts of the country [4-6]. Many of these products are produced using artisanal technologies on-farm and the types and processing steps of these dairy products can vary considerably from one area to the other.

The mainstay of the population in the Amhara region is rain-fed subsistence agriculture. About 73% of the smallholders practice mixed crop-livestock farming, 19% practice crop cultivation, while the remaining 8%

undertake livestock rearing [7]. Like in other regions of Ethiopia, milk production is an integral part of the farming system. Local milk production is mainly from indigenous Zebu cattle which are kept by about half a million smallholder farming households [8] most of whom are poor. In Gondar town dairy production play a significant role in the household income as well as home consumption and also the demand of milk and milk products increased from time to time in the area due to increasing human population, urbanization and other factors. So that identification and understanding of traditional dairy products, handling, processing and utilization of milk and milk products were essential in order to devise appropriate development interventions that would result in improved production and quality of dairy products. Therefore, the overall aim of this study was to assess the practices of handling, processing, and utilization of milk and milk products in Gondar town.

MATERIALS AND METHODS

Description of the Study Area

The study was conducted from November to May, 2016 in Gondar town which was found in Amhara Region state in North Western Ethiopia. The area is located at a distance of 737 km north of Addis Ababa. The area lies between an altitude of 12°35'60"N and longitude of 37°28' 20"E and has an elevation of 2300 meters above sea level (masl). Gondar has a varied landscape, dominantly covered with ragged hills and plateau formations. The annual average temperature was 19.7°C and its annual rainfall was 1772 mm. It could be categorized under woyna-dega climatic zone. The area is also classified mainly in to two seasons, the wet season, from June to September and the dry season from October to May. [9].




Figure 1. Map of study the area.

Data collection and sampling techniques

Data to be collected from the three kebeles namely Kebele 18, 19, and 20 were employed by using purposive sampling techniques. A total of 45 respondents were purposively selected. Both primary and secondary data sources were used for the study. Primary data sources included the household heads and dairy cooperatives in the respective districts. The secondary data was taken from zonal ARDO, NGO operating and from documents that have been written about the study area.

Statistical Analysis

The data was subjected to statistical analysis using statistical package for social sciences (SPSS) software, version 16.0. Descriptive statistics such as frequency, distribution and percentages were used.

RESULT AND DISCUATION

Demographic Characteristics of Respondents

Demographic characteristics of respondents in the study area were showed in Table 1. In the study areas majority of the respondents were male 86.7% and only 13.3% were female. Major of the respondents (42.2%) were found at the age between 47- 62, and the remaining 13.3, 35.6 and 13.3% were found at the age group between 15-30, 31-46 and greater than 62 years old, respectively. Their educational level showed that, 46.7% were read and write and others were illiterate (15.6%), attending elementary education (13.33%), secondary school (8.89%), attend above secondary school (11.11%) and spiritual or religious education (4.4%). Their marital status showed that 91.11% were married, 6.7% were divorced and the remaining 2.22% were single.

Table 1. Demographic characteristic of respondents in Gondar town

Gender	N	%
Male	39	86.7
Female	6	13.3
Total	45	100
Age		
15-30	6	13.3
31-46	16	35.6
47-62	19	42.2
>62	6	13.3
Total	45	100
Educational Level		
Illiterate	7	15.6
Read and Write	21	46.7
Elementary	6	13.33
Secondary	4	8.89
Above secondary	5	11.11
Spiritual	2	4.44
Total	45	100
Marital status		
Single	1	2.22
Married	41	91.11
Divorced	3	6.7
Total	45	100

Family Size

The average family size and percentage of male and female in the study area were shown in Figure 2 and 3, respectively. Based on the information obtained from the respondents the overall average family size in the study area was 5.7. From the total of the population male and female population showed that 47.85% were females and 52.15% were males. From this result it could be conclude that males are more engaged in the dairy production than females.




Figure 2. Average family size in the study area




Figure 3. The percentage of male and female in the study area

Cattle Population in the study area

The overall cattle population in the study area was shown in Table 2. Based on the information obtained from the respondents the overall cattle population in the study area was 419, of which 55 local cows, 138 crossbred cows, 16 local heifers, 27 crossbred heifers, 37 local bulls, 26 crossbred bulls, 33 local calves and 87 crossbred calves.

According to the report of the present study the main milk sources of the area were gain from cross breed dairy cows. However, the study was in agreement with that of the finding of Yilma et al. [10] which stated that milk production depends mainly on indigenous livestock genetic resources especially true in developing countries where production dominated by small holder farmer; more takes place under cattle. The indigenous breeds accounted for 99.19 percent, while the hybrids and pure practices. Thus, farmers and all who handle milk before exotic breeds were represented by 0.72 and 0.09 percent, respectively.

Table 2. The overall cattle population in Gondar town

Kebeles	Number of Cattle Population							
	Cows		Heifers		Bulls		Calves	
	Local	Cross	Local	Cross	Local	Cross	Local	Cross
Kebele 18	18	69	5	8	25	3	14	33
Kebele 19	15	39	3	8	4	7	11	31
Kebele 20	22	30	8	11	8	13	8	23
Total	55	138	16	27	37	26	33	87
Average	18.33	46	5.33	9	12.33	8.66	11	29

Equipment used for milking and storage in the study area

Equipment used for milking and storage were shown in Figure 4. In the study area respondents were used different milking materials for milking and storage of milk. Majority of them were used plastic jar which was 64.4% and 28.9% use gourd and 6.67% of them were used aluminum jar.

The result was agreed with the study of [5, 6] the use of gourd as a churn and a storage vessel. However, it was in agreed with the finding of O'Mahony et al. [11]. In general, the use of clay pot for storage of various dairy products and its use for churning observed.




Figure 4. Equipment used for milking and storage.

Milking and handling of milk and its products

Milking and handling of milk and its product in Gondar town were shown in Table 3. All respondents were washing their hand before milking the caws and 62.2% were washing the udder of the caw before milking and 37.8% were not wash the udder's of cow. After washing, drying of udder was highly recommended. But 66.7% of them were not drying of the udder before milking. In the study area, 55.6% was not properly handled milk and its products in the refrigerator which was the main reason for loss and wastage of milk and its products in the area. However, the remaining 44.4% of respondents were used refrigerator for milk and milk product.

Milk processing materials used in the study area

Milk processing materials in Gondar town were shown in Table 4. In the area respondents were processing milk both in traditional and modern ways. Major of respondents 48.89% were used traditional material i.e gourd and 11.11% were used clay pot whereas some 15.5% were processing milk by using cream separator. The

remaining 24.5% were used other material for processing of milk. The present study was agreed with the study of [5, 6] gourd was the major milk processing container.

Milk processing practice in the Study Area

Milk processing practice in Gondar town was shown in Table 5. Majority of the producers (60%) were follow traditional method of milk processing, and 17.78% were using modern processing materials and the remaining 22.2% were no further processed the milk.

The present study was in line with the report of Duguma and Janssens [15] in Jimma town the traditional milk processing materials were used and methods used in the study area were time consuming, laborious and inefficient in terms of fat recovery.

Table 3. Milking and handling of milk and its product in Gondar town

Variable	N	%
Wash hand	100	100
wash udder	27.99	62
not wash udder	17.01	37.8
Total	45	100
Dry udder	29.7	66.70
Not dry udder	15.3	33.70
Total	45	100
Use refrigerator	19.98	44.40
With no refrigerator	25.02	55.6
Total	45	100

Table 4. Milk processing materials in Gondar town

Milking Processing Equipment	N	%
Cream Separator	7	15.5
Gourd	22	48.89
clay pot	5	11.11
other	11	24.5

Table 5. Milk processing practice in Gondar town

Processing	N	%
Traditional	27	60.00%
Modern	8	17.78%
no process	10	22.2%
Total	45	100%

Preferable Milk processed products in the Study Area

Preferable Milk processed products in Gondar town were shown in Table 6. In the study area skim milk, butter, cheese, whey and yogurt were the preferred milk processed products. In the study area skim milk and butter were more required by respondent. From the respondents 48.89 and 40% were highly preferring milk and butter for household consumption as wells for marketing purposes respectively.

The processed products of the present study was in line with the report of Duguma and Janssens [15] the major products of the traditional milk processing were naturally fermented milk, traditional butter, butter milk, cottage cheese, whey and ghee.

Milk utilization

Milk utilization in Gondar town is indicated in Table 7. Respondent were used different milk product for house hold consumption such as yogurt, milk, cheese, whey. The majority of respondents (51.11%) were preferred to use yogurt for consumption.

The result was in line with the finding of Yilma et al. [13]. Ergo is one of the most common traditionally made fermented milk products in Ethiopia. The result also agreed with Duguma and Janssens [15] the majority of the respondents were process milk into sour whole milk (ergo), cottage cheese (Ayib), butter and ghee.

Table 6. Preferable Milk processed product in Gondar town

Variable	N	%
Skim milk	23	48.89
Butter	18	40
Cheese	2	4.44
Whey	0	0
Yogurt	3	6.67

Table 7. Milk utilization in Gondar town

Variables	N	%
Yogurt	23	51.11
Milk	5	11.11
yogurt and milk	8	17.77
Cheese	1	2.22
Whey	2	4.44
All	6	13.33
Total	45	100

Marketing of milk and milk products

Milk product used for marketing is indicated in Table 8. In the study area milk producers were selling milk to consumers. major of respondents 77.77% were selling milk to consumer, 11.11% were selling milk and butter, 6.66% were by converting milk into butter and they sell butter, the remain 4.44% were selling all milk product to the society.

The study was not agreed with the reports of Beyene [4] in the southern region, Yilma et al.[13] in the central highlands of Ethiopia, Tola [5] in eastern Wollega and Fita [6] in the east Shoa zone of the Oromia region, where most of the farmers do not sell fresh milk but sell butter. On the other hands this result were in line with the finding of Belachew et al. [14] which stated that the producers deliver milk to consumers or consumers may collect it at the producer's gate. Studies also indicated that in terms of volume 71% of intra-urban producers sell milk directly to consumers.

Table 8. Milk product used for marketing

Variable	N	%
Butter	3	6.66
milk and butter	5	11.11
milk	35	77.77
All	2	4.44
total	45	100

CONCLUSION AND RECCOMENDATION

Generally the overall results of the present study indicated that the handling, processing and utilization of milk and its product in Gondar town were lay in traditional and modern way. Milk was processed into different products such as butter, cheese, whey and yogurt. Milk processing was important for household consumption and marketing. In this area producers were consume yogurt in majority. Whereas milk were major product that were highly required for market. Handling of milk and its products were largely in traditional way. Generally in this area large number of respondents was do not have milk and its product processing material adequately, such as refrigerator and other modern material were not accused. Based on the above conclusion the following recommendations were forwarded; government should be try to fulfill the requirement of electric power in different rural kebeles for the use of refrigerator, the government should be offered extension service for the dairy producers for quality of milk, the producers should be used modern milk processing machinery rather

processing in a traditional way, milk collection and processing center should additionally establish and milk pasteurizing and packaging training should be given to the milk produce.

Competing interests

The authors declare that they have no competing interests.

REFERENCES

1. Getachew, F. 2003. Milk and Dairy Product, Post harvest Loss and Food Safety in Sub Saharan Africa and the Near East. A Review of the Small Scale Dairy Sector –Ethiopia. FAO Prevention of Food Losses Programme. FAO, Rome, Italy.
2. Asrat F. 2010. Production, Utilization and Losses Programme. rome: rome publisher.
3. Yilma, Z., Emmanuelle G. 2011. Review of the Ethiopian Dairy Sector. addis abeba, Amhara. Khair, O.E. Elbushra.
4. Beyne, F. 1994. Present situation and future aspects of milk production, milk handling and processing of dairy products in Southern Ethiopia. PhD Thesis, Agricultural University of Norway. Norway.
5. Tola, A. 2002. Traditional milk and milk products handling practices and raw milk quality in Eastern Wollega. MSc Thesis, Alemaya University. Ethiopia.
6. Fita, L. 2004. Assessment of butter quality and butter making efficiency of new churns compared to smallholders' butter making techniques in East Shoa zone of Oromia, Ethiopia. MSc Thesis, Alemaya University. Ethiopia.
7. ZDA, 2005. Annual progress report. West Gojjam Zone Department of Agriculture, Ethiopia
8. CSA, 2009. Agricultural sample survey 2008/09. 16. Report on livestock and livestock characteristics.
9. Gondar, 2008. Gondar Agricultural Rural Development Office. Gondar, Ethiopia.
10. Yilma, Z. Emmanuelle G. and Ameha S. 2011. A Town, Wolaita, South Ethiopia. M.Sc. Thesis.
11. O'Mahony, F. and Bekele E. 1985. Traditional butter making in Ethiopia and possible improvements. ILCA Bulletin 22. International Livestock Center for Africa (ILCA), Ethiopia, Addis Ababa. pp: 9-14.
12. Asamenew Tassew, 2014. Small-Scale Milk Processing, Utilization and Marketing of Traditional Dairy Products In Bahir Dar Zuria And Mecha Districts, Northwestern Ethiopia.
13. Yilma, Z., GuerneBleich, E., Desta, H. & Mugisha, A. 2011. Innovations, Actors and Linkages in the Ethiopian Dairy Value Chain. Proceedings of a National Workshop on Research-Farmer-Extension Linkage in the Dairy Value Chain of Ethiopia held on 29th May 2010. Food and Agriculture Organization of the United Nations, Sub-Regional Office for Eastern Africa, Addis Ababa, Ethiopia. 34-56 p.
14. Belachew Hurisa, Ahmed Mohammed, Haileleul Teferi and AbebeLema. 1994. Dairy products marketing survey in Addis Ababa and the surroundings. DDE (Dairy Development Enterprise), Addis Ababa, Ethiopia.
15. Duguma B and Janssens G.P.J. 2014. Smallholder Milk Processing and Marketing Characteristics at Urban Dairy Farms in Jimma Town of Oromia Regional State, Ethiopia.

Phenotypic Characterization of Indigenous Cattle Populations in West Gojjam Administrative Zones, Amhara National Regional State, Ethiopia

Andualem Tenagne^{1✉}, Getinet Mekuriaw² and Dillip Kumar¹

¹Bahir Dar University, College of Agriculture and Environmental Sciences, P.O. Box 79, Bahir Dar, Ethiopia

²International Livestock Research institute (ILRI) P.O. Box 5689, Ethiopia Addis Abeba, Ethiopia

✉Corresponding author's Email: andu2026@gmail.com

Received 28 Sep. 2016 Accepted 05 Nov. 2016

ABSTRACT: The study was carried out in Semen Achefer, Sekela and Jabitenan districts of Western Gojjam zone of Amhara National Regional State. The objective of the study was to carry out phenotypic characterization of local cattle population in the study under farmers' management condition in the study area. A total of 600 cattle were sampled randomly for characterization of phenotypic traits. Data were gathered through field observations and linear body measurements of sample populations. The Sampled indigenous cattle were identified by sex and district (agro ecology). The most dominant coat colour patterns in the sampled populations were plain, patchy and spotted with the most frequently observed coat colour type being light red, black and dark red. Sex of animals had $P < 0.05$, on all of the body measurements. Agro ecology also showed $p < 0.05$, for most of the body measurements, except tail length, horn length, height at wither and rump height. Among leaner body measurements moderate correlations and positive relationship were shown. The prediction of body weight could be based on regression equation $y = -481.55 + 4.89x$ for male sample population and $y = -405.22 + 4.64x$ for female sample cattle population where y and x are body weight and chest girth, respectively. Most of the body measurements of cattle were affected by sex and agro- ecology. Phenotypic result of cattle populations in the study areas was varied from former finding and therefore; to put specific characteristics' of the breed, further molecular characterization is needed.

Author Keywords: Body Weight, Cattle, Characterization, Indigenous, Linear Body Measurement

Abbreviations: CSA: Central Statistical Agency; DA: Development Agent; DAGRIS: Domestic Animal Genetic Resource Information System; FAO: Food and Agricultural Organization; GLM: General Linear Model; LBM: Linear Body Measurements Masl: MetereAbove Sea Level; Mm: Milimetre; °C Degree centigrade; WAO: Woreda Agricultural Office

INTRODUCTION

Ethiopia has served as a gateway to domestic animals from Asia to Africa and its diverse ecology favored diversification of these resources [1]. The country is endowed with huge livestock resources of varied and diversified genetic pools with specific adaptations to a wide range of agro-ecologies [2, 3]. Among livestock species, cattle have significant contributions to the livelihoods of the farmers. They serve as a source of draught power for the rural farming population, supply farm families with milk, meat, manure, serve as source of cash income, and play significant role in the social and cultural values of the society. Cattle contribute nearly all the draught power for agricultural production at smallholder level in Ethiopia [4].

The total number of cattle in all regions of the country except the non-sedentary population of three zones of Afar and six zones of Somali region was estimated to be 57 million, has the largest population in Africa [5]. The majority of these cattle (98.95 percent) are indigenous breeds which are kept under extensive management [1]. This is because indigenous cattle have been naturally selected for years towards adaptive traits as tolerance and resistance to diseases, high fertility, unique product qualities, longevity and adaptation to harsh environments and poor quality feeds [6]. However, a large proportion of indigenous livestock populations in the developing world have not yet been characterized or evaluated at phenotypic and genetic levels [7]. In order to ensure proper conservation and utilization of indigenous breeds, it is necessary to evaluate genetic variations that exist within

and among breeds. Accordingly, proper identification, evaluation, and maintenance of different traits of animal genetic resources are necessary to make them available and relevant for future use without compromising their current utilization [8]. Phenotypic as well as genetic characterization of indigenous livestock genetic resources provides the basis for any livestock development intervention. Clearly, sustainable utilization of local breeds is the best means of conserving these genetic resources. The first essential step towards sustainable utilization of these resources is to identify the major breed types, establish their population size as well as their geographical distribution and describe their typical qualitative and quantitative phenotypic traits [9]. Recognition of breeds' potential depends on the availability of accurate and comprehensive information on their characteristics and their production and marketing environments. Such information can only be obtained through well-designed characterization studies that include pertinent and well thought-out analysis and interpretation of the data collected [10]. However, the genetic diversity and the genetic merits of most Ethiopian indigenous cattle populations are not yet well understood and exploited. Consequently, some of the indigenous cattle populations are already extinct and endangered, while the risk status of many of them is unknown [11]. Despite the significant contribution of cattle to the country, little attention is given to identify, characterize and conserve the diversity of the various classes of livestock. The current state of knowledge on characterization of cattle genetic resources in Ethiopia shows that there is inadequate breed level characterization information [12].

West Gojjam zone is one of the administration zone Amhara National regional State which has high cattle population potential and suitable weather conditions for cattle production. Even though the area has suitable environment and great potential for cattle production, there was a gap in utilizing its maximum potential and proper conservation and utilization of indigenous cattle breeds. Therefore in the study area; there was a need to conduct phenotypic characterization to solve the existing problems in the area. The objective of the study was to characterize phenotypic characteristics of local cattle population in the study area

MATERIALS AND METHODS

The study was conducted in 3 districts of West Gojjam zone of the Amhara National Regional State, Ethiopia. West Gojjamzone is one of administrative zone found in Amhara regional state and which is located on the southern border of Lake Tana. The districts included in the study sites were North Achefer, Sekela and Jabitenan (Figure 1).




Figure 1. Map of the study area (Ethiopia; Amhara National Regional State; study districts)

Sampling Technique and Procedure

Multi-stage purposive sampling technique was employed to select the districts and kebeles for the study. Study districts were stratified based on agro ecology in to three strata; lowland, midland and highland. Three kebeles were selected from each stratum purposively based on cattle population potential and agro ecology (Table 1). Finally, 30 households (cattle owners) that have more than 2 head of cattle for interview and 67 animals (cattle) for measurement were selected randomly from each kebele. The sample size was calculated based on [13].

Table 1. Sampling Frame of Study areas

Districts	Agro-ecology of districts	Sample kebeles	Sample House holds	Sample Animals	Altitude Masl & Rain Fall
Semen Achefer	M.L	0	0	0	
	L.L	3	90	200	<1500&<800mm
Sekela	H.L	3	90	200	>2500&1200-2200mm
	M.L	0	0	0	
Jabitenan	M.L	3	90	200	1500-2500&800-1200mm
	L.L	0	0	0	
Total		9	270	600	

M.L= mid land, L.L = low land, H.L= high land

Data Collection

General information of the area, topography, climatic data and population size were obtained from secondary data from districts agricultural development offices. In each sampling site, the selected cattle owners were briefed about the importance and objectives of the study before the commencement of the actual data collection. Visual observation was made and morphological features were recorded based on breed morphological characteristics descriptor list [14, 15]. Linear body measurements were taken using a standard textile measuring tape and standard steel tape. Qualitative and quantitative traits were recorded through prepared check list from 192 mature males and 408 mature females. A total of 12 qualitative traits were examined and recorded: body hair color pattern, body hair coat color, udder size, horn presence, horn shape, horn orientation, ear orientation, hump size, navel flap (for cows), preputial sheath (for bulls), facial (head) profile and tail length.

A total of 11 quantitative traits were measured and recorded: heart girth, body length, height at withers, height at rump, pelvic width, ear length, horn length, cannon bone circumference, mouth circumference and body weight. The morphological variables recorded in this study were adapted from the standard cattle breed descriptor list [7] and extensively used in Ethiopia [15- 17]. Every animal to be measured was identified by sex and study site.

Data Management and Analysis

Qualitative and quantitative body measurement data were first entered into Excel 2007 computer software and analyzed using SPSS version 20. Qualitative data were analyzed using the frequency procedure of SPSS. While quantitative data were analyzed using the Generalized Linear Model (GLM) procedure of SAS. Sex and district (agro ecology) were fitted as fixed effects while linear body measurements were fitted as dependent variables. When analysis of variance declares significance, least square means were separated. Pearson's correlation coefficients were estimated among body weight and linear body measurements and between linear body measurements for females and males (SPSS version 20). Correlations (Pearson's correlation coefficients) between body weight and the linear measurements were computed for the population within each sex. To quantify the effect of independent variables (site and sex) on the linear body measurement (dependent variables) of the sample populations, the GLM procedure of SPSS 20 was employed. The model fitted for linear body measurements for sample populations was, $Y_{ijk} = \mu + A_i + S_j + e_{ijk}$

Where,

Y_{ijk} = Observed value of the trait of interest

μ = Overall mean

A_i = Fixed effect of ith agro ecology (sites)

S_j = Fixed effect of jth sex

e_{ijk} = Residual random effect

RESULTS AND DISCUSSION

Phenotypic Characterization

On farm phenotypic characterization of cattle breed includes all the qualitative description and morphological measurements of the animal. It is a primary and low cost animal genetic resource characterization as compared to the on-station [18].

Qualitative trait of indigenous cattle types found in Low land, High land and Mid land agro-ecologies are presented in Table 2. The most frequent color patterns observed in the study area were plain 63.17%, patchy 18.5% and spotted 18.33%. Out of the 63.17% (largest), plain coat color pattern, 39.67% light red, 21.33% black, 17% dark red, 14.33% grey and 7.67% fawn were the dominant color types (Figure 2). Comparably in Fentalle district the dominantly observed coat colour patterns for Kereyu cattle were 31.7%, 33.3% and 35 % for plain, patchy and spotty, respectively [19]. And in mursi areas frequently observed coat patterns were plain 52.0%, pied 36.0% and spotted 12.0% [20]. Light red coat colour was mostly observed in the study area Table 2 Similar to Raya Sanga [16]. In contrast to this, for Boran cattle white was dominant coat color [8], white and black frequently observed coat colour for Mursicattle breeds [20] and Grey color was the most observed forkereyu cattle breeds [19].

In the study area among the sampled cattle population the majority (96.83%) of cattle had horn, whereas, 3.17% were polled. Out of 96.83% horned cattle population 58.5% straight, 36.67% curved and 4.17% lyre were mainly observed horn shapes in the study areas. 47.5% of horns were oriented tips pointing laterally, 25.5% upwards, 25.5% forward and 1.5% down wards (Figure 3). Hump size difference were observed between male and female population in each district (Table 2 and 3).



Figure 2. Different coat colour of local cattle in Mid and Low land agro-ecologies respectively.



Figure 3. Hump size and horn orientation of local cattle in the study areas

Table 2. Qualitative trait description of male animal in West Gojjam zone

Districts (agro ecology) Phenotypic variables	Low land	High land	Mid land	Overall mean
	N(%)	N(%)	N(%)	N(%)
Coat color Patter				
➤ Plain	45(65.2)	26(50.0)	44(62.0)	115(59.9)
➤ Patchy	14(20.3)	12(23.1)	13(18.3)	39(20.3)
➤ Spotted	10(14.5)	14(26.9)	14(19.7)	38(19.8)
	$X^2\text{value} = 60.97^*$			
Hair Coat color				
➤ Black	14(20.3)	8(15.4)	16(22.5)	38(19.8)
➤ Dark red	7(10.1)	11(21.2)	9(12.7)	27(14.0)
➤ Light red	36(52.2)	21(40.4)	27(38.0)	84(43.8)
➤ Fawn	3(4.3)	5(9.6)	6(8.5)	14(7.3)
➤ Grey	9(13.0)	7(13.5)	13(18.3)	29(15.1)
	$X^2\text{value} = 75.3^*$			
Horn presence				
➤ Horned	67(97.1)	49(94.2)	69(97.2)	185(96.4)
➤ Polled	2(2.9)	3(5.8)	2(2.8)	7(3.6)
	$X^2\text{value} = 165.0^*$			
Horn shape				
➤ Straight	47(68.1)	29(55.8)	41(57.7)	117(60.9)
➤ Curved	20(29.0)	20(38.5)	24(33.8)	64(33.3)
➤ Lyre-shape	1(1.4)	2(3.8)	6(8.9)	9(4.7)
➤ Loose	1(1.4)	1(1.9)	0	2(1)
	$X^2\text{value} = 180.3^*$			
Horn orientation				
➤ lateral	30(43.5)	21(40.4)	40(56.3)	91(47.4)
➤ Upward	22(31.9)	13(25.0)	16(22.5)	51(26.6)
➤ Downward	1(1.4)	0(0)	2(2.8)	3(1.6)
➤ Forward	16(23.2)	18(34.6)	13(18.3)	47(24.5)
	$X^2\text{value} = 80.9^*$			
Ear orientation				
➤ Erect	0	1(1.9)	0	1(0.05)
➤ Lateral	67(97.1)	51(98.1)	70(98.6)	188(97.9)
➤ Drooping	2(2.9)	0	1(1.4)	3(1.6)
	$X^2\text{value} = 360.3^*$			
Hump size				
➤ Small	11(15.9)	4(7.7)	14(19.7)	29(15.1)
➤ Medium	43(62.3)	17(32.7)	30(42.3)	90(46.9)
➤ Large	15(21.7)	29(55.8)	27(38.0)	61(31.8)
	$X^2\text{value} = 99.4^*$			
Perpetual Sheath (bull)				
➤ Small	14(20.3)	12(23)	8(11.3)	34(8.3)
➤ Medium	31(44.9)	26(50.0)	37(52.1)	94(48.9)
➤ Large	24(34.8)	14(26.9)	26(36.6)	64(15.7)
	$X^2\text{value} = 119.4^*$			
Facial (head)				
➤ Straight	60(87.0)	48(92.3)	63(88.7)	1171(89)
➤ Concave	3(4.3)	3(5.8)	3(4.2)	9(4.7)
➤ Convex	6(8.7)	1(1.9)	5(7.0)	12(6.3)
	$X^2\text{value} = 268.4^*$			
Tail length				
➤ Short	1(1.4)	0	1(1.4)	2(0.1)
➤ Medium	2(2.9)	0	1(1.4)	3(1.6)
➤ Long	66(95.7)	52(100)	69(97.2)	187(97.4)

Pr. = perpetual, facial = facial profile

Table 3. Qualitative trait description of female animal in West Gojjam zone

Districts (agro ecology)	Low land	High land	Mid land	Overall mean
Phenotypic variables	N(%)	N(%)	N(%)	N(%)
Coat color Patter				
➤ Plain	84(64.1)	87(58.8)	93(72.1)	264(64.7)
➤ Patchy	21(16.0)	28(18.9)	23(17.8)	72(17.96)
➤ Spotted	26(19.8)	33(22.3)	13(10.1)	97(23.8)
	$X^2\text{value} = 179.3^*$			
Hair Coat color				
➤ Black	33(25.2)	32(21.6)	25(19.4)	90(22.0)
➤ Dark red	15(11.5)	38(25.7)	22(17.1)	75(18.4)
➤ Light red	60(45.8)	41(27.7)	53(41.1)	154(37.8)
➤ Fawn	5(3.8)	23(15.5)	4(3.1)	32(7.8)
➤ Grey	18(13.7)	14(9.5)	25(19.4)	57(13.9)
	$X^2\text{value} = 103.2^*$			
Horn presence				
➤ Horned	128(97.7)	141(95.3)	127(98.4)	396(97.1)
➤ Polled	3(2.3)	7(4.7)	2(1.6)	12(2.9)
	$X^2\text{value} = 165.0^*$			
Horn shape				
➤ Straight	79(60.3)	83(56.1)	72(55.8)	234(57.4)
➤ Curved	40(30.5)	60(40.5)	56(43.4)	156(38.2)
➤ Lyre-shape	11(8.4)	4(2.7)	1(0.8)	16(8.3)
➤ Loose	1(0.8)	1(0.7)	0	2(0.5)
	$X^2\text{value} = 365.6^*$			
Horn orientation				
➤ lateral	66(50.4)	65(43.9)	63(48.8)	194(47.5)
➤ Upward	26(19.8)	42(28.4)	34(26.4)	102(25)
➤ Downward	2(1.5)	0	4(3.1)	6(1.5)
➤ Forward	37(28.2)	41(27.7)	28(21.7)	106(26)
	$X^2\text{value} = 171.3^*$			
Ear orientation				
➤ Erect	0	7(4.7)	0	7(1.7)
➤ Lateral	130(99.2)	141(95.3)	119(92.2)	390(95.6)
➤ Drooping	1(0.8)	0	10(7.8)	11(2.7)
	$X^2\text{value} = 708.0^*$			
Uder Size				
➤ Small	26(19.8)	27(18.2)	26(20.2)	79(19.4)
➤ Medium	64(48.9)	85(57.4)	59(45.7)	208(50.9)
➤ Large	41(31.3)	36(24.3)	44(34.1)	121(29.7)
	$X^2\text{value} = 212.6^*$			
Hump size				
➤ Small	122(93.1)	109(73.6)	125(96.)	356(87.3)
➤ Medium	9(6.9)	35(23.6)	4(3.1)	48(11.8)
➤ Large	0	4(2.7)	0	4(0.9)
	$X^2\text{value} = 537.6^*$			
Navel flap				
➤ Small	29(22.1)	46(31.1)	24(18.6)	99(24.6)
➤ Medium	75(57.3)	72(48.6)	71(55.0)	218(53.4)
➤ Large	27(20.6)	20(13.5)	34(26.4)	81(19.9)
	$X^2\text{value} = 624.9$			
Facial (head)				
➤ Straight	114(87)	143(96.6)	117(90.7)	374(91.7)
➤ Concave	10(7.6)	2(1.4)	9(7.0)	21(5.1)
➤ Convex	7(5.3)	3(2.0)	3(2.3)	13(2.0)
	$X^2\text{value} = 629.6^*$			
Tail length				
➤ Short	2(1.5)	0	1(0.8)	3(0.07)
➤ Medium	2(1.5)	12(8.1)	4(3.1)	18(4.4)
➤ Long	127(96.9)	136(91.9)	124(96.1)	390(95.6)
	$X^2\text{value} = 697.7^*$			

Hump size of male cattle population were 15.9% small, 62.3% medium and 21.7% large in Low land, 7.7% small, 32.7% medium and 55.8% large in High land and 19.7% small, 42.3% medium and 38% large in Mid land agro-ecology. Whereas, for cow 93.1% small and 6.9% medium in Low land, 73.6% small, 23.6% medium and 2.7% large High land and 96.9% small and 3.1% medium in Mid land agro-ecologies.

The overall facial profile of the three districts was 90.83% straight 5% concave and 4.17% convex. Likewise, their ear orientations were 1.3% erected, 96.33% laterally oriented and 2.3% dropping oriented. A total of both male and female cattle population in the study areas having tail length of 95.67% long (below the hocks), 3.5% medium (about the hocks) and 0.83% short (above the hocks) (Figure 4). The perpetual sheath of male sample population were 20.3% small, 44.9% medium and 34.8% large in Low land, 23.1% small, 50% medium and 26.9% large in High land and 11.3% small, 52.1% medium and 36.6% in Low land agro-ecology.

From the total female cattle population evaluated, udder size of 19.8% them were small, 48.9% medium and 31.3% large in Low land, 18.2% small, 57.4% medium and 24.3% large in High land and 20.2% small, 45.7% medium and 34.1% large in Mid land agro-ecology (Figure 5). Similarly navel flap for cows were 22.1% small, 57.3% medium and 20.6% large in Low land, 31.1% small, 48.6% medium and 13.5% large in High land and 18.6% small, 55% medium and 26.4% large in Mid land agro-ecologies. As shown in Table 2, all of qualitative traits were significantly different among districts and this difference might be due to the agro ecological difference of the three districts.



Figure 4. Perpetual sheath and tail length of the bull in Mid land and Low land agro-ecology respectively.



Figure 5. Udder size and naval flap of cow respectively in High land agro-ecology

Table 4. Body measurements (cm) of adult local cattle population in the study area (LSM±SE)

Variables	MC		HL		EL		HW		BL	
	N	LSM±SE	N	LSM±SE	N	LSM±SE	N	LSM±SE	N	LSM±SE
Over all	600	38.9±0.07	600	22.84±0.42	600	19.83±0.06	600	114.5±0.24	600	115.41±0.24
Agro-ecology	--	*		NS		*		NS		*
Low land	200	38.3±0.12	200	22.4±0.66	200	19.4±0.09	200	115±0.4	200	116.7±0.3
High land	200	39.2±0.12	200	22.6±0.69	200	19.8±0.1	200	114±0.4	200	114.4±0.3
Mid land	200	39.2±0.12	200	23.5±0.67	200	20.2±0.09	200	114.5±0.4	200	115±0.3
Sex	--	*		*		*		*		*
Male	192	40.07±0.12	192	23.9±0.7	192	20.2±0.1	192	116.2±0.4	192	117.06±0.3
female	408	38.8±0.08	408	21.8±0.5	408	19.5±0.07	408	112.8±0.3	408	113.7±0.2

NS= Non significant; N = number of household; LMS =least square mean; SE = standard error

Table 4. (Continued)

Variables	HG		PW		TL		RH		CBC		BW
	N	LSM±SE	N	LSM±SE	N	LSM±SE	N	LSM±SE	N	LSM±SE	N(LSM±SE)
Over all	600	146.04±0.27	600	36.5±0.09	600	78.6±0.22	600	115.8±0.25	600	20.3±0.06	600 (266.8±2.3)
Agro- ecology	--	*		*		NS		NS		*	*
Low land	200	144.8±0.43	200	36.3±0.14	200	78.8±0.35	200	115.8±0.42	200	20.3±0.09	200 (260±2.18)
High land	200	147.2±0.4	200	36.4±0.15	200	79±0.4	200	115.8±0.43	200	20±0.09	200 (272.2±2.4)
Mid land	200	146.±0.44	200	37±0.14	200	78.1±0.35	200	115.9±0.4	200	20.5±0.09	200 (268.3±2.2)
Sex	--	*		*		*		*		*	*
Male	192	151.8±0.44	192	37.08±0.15	192	79.6±0.36	192	117.2±0.14	192	21.14±0.09	192 (300.7±4.3)
female	408	140.3±0.30	408	35.98±0.1	408	77.7±0.24	408	114.5±28	408	19.5±0.06	408 (243.6±2.3)

NS= Non significant; N = number of household; LMS =least square mean; SE = standard error

Table 5. Coefficient of correlations between body weight and linear body measurements (Above diagonal for male and below diagonal for female)

	MC	HL	EL	HW	BL	HG	PW	TL	RH	CBC	BW
MC		0.210**	0.304**	0.479**	0.259**	0.566**	0.309**	0.368**	0.293**	0.567**	0.564**
HL	0.215**		0.002 ^{ns}	0.212**	0.332**	0.116 ^{ns}	0.100 ^{ns}	0.163*	0.117 ^{ns}	0.409**	0.123 ^{ns}
EL	0.440**	0.047		0.187**	0.169 ^{ns}	0.406**	0.377**	0.316**	0.125 ^{ns}	0.318**	0.393**
HW	0.198**	0.101**	0.117**		0.530**	0.431**	0.290**	0.337**	0.537**	0.462**	0.439**
BL	0.310**	0.172**	0.171**	0.276**		0.330**	0.303**	0.283**	0.303**	0.415**	0.339**
HG	0.452**	0.057 ^{ns}	0.244**	0.275**	0.385**		0.547**	0.497**	0.324**	0.510**	0.984**
PW	0.393**	0.136**	0.482**	0.187**	0.238**	0.301**		0.370**	0.245**	0.368**	0.547**
TL	0.366**	0.215**	0.268**	0.278**	0.314**	0.477**	0.355**		0.324**	0.382**	0.496**
RH	0.386**	0.176**	0.162**	0.588**	0.480**	0.509**	0.222**	0.436**		0.368**	0.322**
CBC	0.413**	0.069 ^{ns}	0.282**	0.285**	0.445**	0.430**	0.278**	0.304**	0.397**		0.504**
BW	0.439**	0.066 ^{ns}	0.235**	0.278**	0.379**	0.994**	0.293**	0.467**	0.507**	0.423**	

^{ns} non- significance; MC= Mouth Circumference, HL= Horn Length, EL= Ear Length, HW= Height at Withers, RH= Rump Height, BL= Body Length, HG= Heart Girth, PW= Pelvic Width, TL= Tail length, RH= Rump height and CBC= Cannon bone circumference

Morphological measure of adult cattle population

The quantitative measures of some phenotypic traits of local cattle population are summarized in (Table 4). The overall least squares means and standard error values of mouth circumference, horn length, ear length, height at wither, body length, heart girth, pelvic width, tail length, rump height and cannon bone circumference were 38.9 ± 0.07 cm, 22.8 ± 0.42 cm, 19.83 ± 0.06 cm, 114.5 ± 0.2 cm, 115.4 ± 0.2 cm, 146 ± 0.27 cm, 36.5 ± 0.09 cm, 78.6 ± 0.22 cm, 115.8 ± 0.25 cm and 20.3 ± 0.6 cm, respectively. All phenotypic measurements listed on the above were significantly different ($P < 0.05$) between male and female cattle. Most body measurements of cattle like heart girth, body length, pelvic width, ear length, cannon bone circumference, mouth circumference and body weight were also significant difference ($P < 0.05$) among Low Land, High Land and Mid Land locations. The possible reason for the difference may be agro ecological difference among districts. But horn length, tail length, height at wither and rump height were did not shows significant difference ($P < 0.05$) among the three agro ecologies. Body length, chest girth, and wither height of the male cattle were measured to be 117.6 ± 0.3 , 151.8 ± 0.44 , and 116.25 ± 0.4 cm, respectively. These measurements were found lower than body measurements reported on the same sex for Boran cattle breeds [8], Mursi cattle breeds [20] and for Wollocattle [16] except heart girth. But heart girth of this result was aligned with that reported [16] for Wollo cattle breed. Like other local zebu cattle populations, the male and female of this result showed significantly different for most of morphological measurements and all linear measurement male value greater than female's. Different reports revealed that the mean value of on-farm morphological traits measurement on local male and female cattle result that males are usually greater than their counter female groups [21, 22, and 19]. Therefore cattle populations in the study areas were varied with qualitative and quantitative traits from former findings of Fasil and Dereje [17, 16] respectively and there was no sufficient evidence to classify either of them.

Correlation between Body Weight and Linear Body Measurements

Pearson's correlation coefficient between body weight and linear body measurements for male and female were calculated and presented in the (Table 5). In males all of linear body measurements have positive association with body weight and strong association was found between body weight and chest girth ($r = 0.984$), whereas mouth circumference ($r = 0.56$), tail length ($r = 0.5$), pelvic width ($r = 0.55$) and cannon bone circumference ($r = 0.5$) had moderate correlation with body weight. Wither height ($r = 0.44$), body length ($r = 0.34$), rump height ($r = 0.32$) and ear length ($r = 0.39$) showed mild correlation with body weight. But horn length for both sex male ($r = 0.12$) and female ($r = 0.002$) did not showed significant correlation with body weight. These linear body measurements that showed moderate and mild correlation were may not significantly affected by the change in body weight; hence, they are not more important in prediction of live body weight of the animal. In females also all of linear body measurements have positive association with body weight and strong association was found between heart girth and body weight. Rump height showed moderate correlation ($r = 0.50$), with body weight. Height at wither ($r = 0.28$), body length ($r = 0.38$), mouth circumference ($r = 0.44$), tail length ($r = 0.48$), pelvic width ($r = 0.29$) and cannon bone circumference ($r = 0.42$) were showed mild correlation, whereas ear length ($r = 0.24$) had weak correlation with body weight. Generally as the result of correlation showed heart girth (chest girth) was the most important than other linear body measurement for both male and female to estimate body weight.

Estimated Mature Body Weight of the Sample Population by sex

The estimated average mature body weight as used conversion of from heart girth were 300.7 ± 4.3 for male and 243.6 ± 2.3 kg for female with at rang of (214-388 kg) for male and (164-381kg) for female. These Variations were observed among individuals which were compared to other local cattle [19] ranging from 196.9 to 333.6 for females and from 178.1 to 428.2 kg for males. Based on the estimated body weight of the individual animals the following linear regression equations (body weight on heart girth) were developed separately for both sexes.

$$Y_m = -481.55 + 4.89x \text{ for bulls and}$$

$$Y_f = -405.22 + 4.64x \text{ for cows}$$

Where:

Y_f = estimated body weight of mature female cattle (kg)

Y_m = estimated body weight of mature male cattle (kg)

x = heart girth.

CONCLUSIONS AND RECOMMENDATIONS

The most dominant coat colour pattern was plain and frequently observed coat colour type being light red. The majority of cattle were possessed horn with straight shape and tips pointing lateral orientation. Sample population of bulls had medium hump size and perpetual sheath whereas cows possessed medium udder size and naval flap. Sex of animals had significant effect ($P<0.05$) on body weight and all of the body measurements. District (agro ecology) also had significant effect on body weight and all of the body measurements except height at wither, horn length, tail length and rump height ($P<0.05$). Among the body measurements of sample population moderate and significant ($P<0.05$) positive correlation was found. Chest girth was the most important linear body measurement to estimate body weight.

Overall, cattle populations in the study areas were varied from former local cattle bred that were identified in Gojjam and Wollo areas. Therefore further characterization of local cattle in the study area at molecular level should be duly required.

Acknowledgements

We would like to thank Kebele DA's and districts experts for their unreserved assistance in data collection and body measurement

REFERENCES

1. CSA (Central Statistical Agency). 2013. Agricultural Sample Survey 2012/2013, Volume II Report on Conservation of Animal Genetic Resources. Commission on Genetic Resources for food and Agriculture 5th Session, Rome, 28-30 January 2009.
2. Roessler R, Drucker A G, Scarpa R, Markemann A, Lemke U, Thuy LT and Valle Zárate A. 2008. Using choice experiments to assess smallholder farmers' preferences for pig breeding traits. Roles of Livestock in Major Production System)
3. Gemedaduguma, Mirkena T, Haile A, Iñiguez L, Okeyo AM, Tibbo, M, Rischkowsky B, Sölkner J, Wurzinger, M. 2010. Participatory approaches to investigate breeding objectives of livestock keepers. *Livest Res Rural Dev*. Volume 22, Article 64. Retrieved April 5, 2010.
4. Tefera M. 2011. Oxenization Versus Tractorization: Options and Constraints for Ethiopian Farming System Ethiopia. *International Journal of Sustainable Agriculture* 3 (1): 11-20, 2011, ISSN 2079-2107.
5. CSA (Central Statistical Agency). 2015. Agricultural sample survey report on livestock and livestock characteristics. Central statistical agency federal democratic republic of Ethiopia; Addis Ababa. Volume 2: pp 194.
6. Aynalem Haile. 2006. Genetic and economic analysis of Ethiopian Boran cattle and their crosses with Holstein Friesian in central Ethiopia. PhD thesis. National dairy research institute (I.C.A.R) Karnal-132001 (Haryana), India. 197 p.
7. DAGRIS. 2006. Domestic Animal Genetic resource Information system. DAGRIS (eds. J.E.O. Rege, W. Ayalew, E. Getahun, O. Hanotte and T. Dassei). International Livestock Institution. Addis Ababa. Ethiopia.
8. Tekle S, Dessie T and Kebede K. 2011. on-farm characterization of boran cattle breed in the production systems in Dire district of Borana Zone, Oromia Region, Ethiopia.
9. Ayalew W, Getahun E, Tibbo M, Mamo Y and Rege J.E.O. 2004. Current State of Knowledge on Characterisation of Farm Animal Genetic Resources in Ethiopia. pp. 1-21 In: Proceedings of the 11th Annual Conference of the Ethiopian Society of Animal Production (ESAP), 28-30 August 2003. Addis Ababa, Ethiopia.
10. FAO. 2014. Access and Benefit-sharing of Animal Genetic Resources Using the Nagoya Protocol as a Framework for the Conservation and Sustainable Use of Locally Adapted Livestock Breeds by Ilse Koehler-Rollefson.
11. Edea Z, Dadi H, Wook Kim S, Dessie T and Kwan-Suk Kim. 2012. Comparison of SNP Variation and Distribution in Indigenous Ethiopian and Korean Cattle Populations. P: 1-4.
12. Rowlands J, Nieves C, Hanotte O, Ayalew W. 2006. Cattle breed distributions across districts as determined from cluster analysis of phenotypic data collected in the Oromiya region, Ethiopia. In 8th World Congress on Genetics Applied to Livestock Production, August 13-18, 2006, Belo Horizonte, MG, Brasil.
13. Kothari CR. 2004. *Research Methodology: Methods and Techniques*. 2nd ed. New Delhi, Wiley Eastern.

14. FAO. 2012. Characterization and value addition to local breeds and their product in near east north Africa. Food and agriculture organization of the United Nations. Rome.
15. Wuletaw Z. 2004. Survey of indigenous cattle genetic resources, husbandry practices and cattle breeding objectives in North Gondar, Ethiopia. MSc thesis, Alemaya University, Dire Dawa, Ethiopia.
16. Tadesse D. 2005. On-farm phenotypic characterization of cattle genetic resources and their production systems in south and north Wollo Zones of Amhara Region, North Eastern Ethiopia, MSc Thesis, Alemaya University, Dire Dawa, Ethiopia. pp. 125.
17. Getachew F and Ayalew W. 2014. On-Farm Phenotypic Characterization of indigenous Cattle population in Awi, East and West Gojjam Zones of Amhara Region, Ethiopia. Research Journal of Agriculture and Environmental Management. 3(4): 227-237, April, 2014.
18. FAO. 2007. The State of the World's Animal Genetic Resources for Food and Agriculture. Commission on Genetic Resources for Food and Agriculture Organization of the United Nations, Rome.
19. Garoma Sh, 2006. In-situ Phenotypic Characterization of Kereyu cattle type in Fentale district of Oromia Region, Ethiopia. M.Sc. Thesis. School of Graduate Studies. Haramaya University.
20. Terefe E, Dessie T, Haile A, Mulatu W and Mwai O. 2012. Characterization of Mursi Cattle Breed In Its Production Environment, InSalamago Wereda, South-West Ethiopia. M.Sc. Thesis, Alemaya University, Ethiopia.
21. Taye T. 2005. On-Farm Phenotypic Characterization of Sheko Breed of Cattle and Their Habitat in Bench Maji zone, Ethiopia. MSc Thesis. Alemaya University, Ethiopia.
22. Mekuria G. 2005. On- station *ex-situ* characterization of Ogaden cattle breed at Alemaya University. MSc thesis, Alemaya University, Dire Dawa, Ethiopia.

Instructions for Authors

Manuscript as Original Research Paper, Review and Case Reports are invited for rapid peer-review publishing in the *Journal of Life Science and Biomedicine*. Considered subject areas include: Biocontrol, Biochemistry, Biotechnology, Bioengineering, Neurobiology... [view full aims and scope](#)

[JLSB EndNote Style](#)

[Manuscript Template \(.doc\)](#)

[Sample Articles](#)

[Declaration form](#)

[Policies and Publication Ethics](#)

Submission

The manuscript and other correspondence should preferentially be [submit online](#). Please embed all figures and tables in the manuscript to become one single file for submission. Once submission is complete, the system will generate a manuscript ID and will send an email regarding your submission. Meanwhile, the authors can submit or track articles via editors@jlsb.science-line.com ; jlsb.editors@gmail.com. All manuscripts must be checked (by English native speaker) and submitted in English for evaluation (in totally confidential and impartial way).

Supplementary information

The online submission form allows supplementary information to be submitted together with the main manuscript file and covering letter. If you have more than one supplementary files, you can submit the extra ones by email after the initial [submission](#). Author guidelines are specific for each journal. Our Word template can assist you by modifying your page layout, text formatting, headings, title page, image placement, and citations/references such that they agree with the guidelines of journal. If you believe your article is fully edited per journal style, please use our [MS Word template](#) before submission. **Supplementary materials** may include figures, tables, methods, videos, and other materials. They are available online linked to the original published article. Supplementary tables and figures should be labeled with a "S", e.g. "Table S1" and "Figure S1". The maximum file size for supplementary materials is 10MB each. Please keep the files as small possible to avoid the frustrations experienced by readers with downloading large files.

Submission to the Journal is on the understanding that

- 1.The article has not been previously published in any other form and is not under consideration for publication elsewhere;
- 2.All authors have approved the submission and have obtained permission for publish work.
- 3.Researchers have proper regard for conservation and animal welfare considerations. Attention is drawn to the '[Guidelines for the Treatment of Animals in Research and Teaching](#)'. Any possible adverse consequences of the work for populations or individual organisms must be weighed against the possible gains in knowledge and its practical applications. If the approval of an ethics committee is required, please provide the name of the committee and the approval number obtained.

Ethics Committee Approval


Experimental research involving human or animals should have been approved by author's institutional review board or ethics committee. This information can be mentioned in the manuscript including the name of the board/committee that gave the approval. Investigations involving humans will have been performed in accordance with the principles of [Declaration of Helsinki](#). And the use of animals in experiments will have observed the *Interdisciplinary Principles and Guidelines for the Use of Animals in Research, Testing, and Education* by the New York Academy of Sciences, Ad Hoc Animal Research Committee. If the manuscript contains photos or parts of photos of patients, informed consent from each patient should be obtained. Patient's identities and privacy should be carefully protected in the manuscript.

Graphical Abstract

Authors should provide a graphical abstract (a beautifully designed feature figure) to represent the paper aiming to catch the attention and interest of readers. Graphical abstract will be published online in the table of content. The graphical abstract should be colored, and kept within an area of 12 cm (width) x 6 cm (height) or with similar format. Image should have a minimum resolution of 300 dpi and line art 1200dpi.

Note: Height of the image should be no more than the width.

Please avoid putting too much information into the graphical abstract as it occupies only a small space. Authors can provide the graphical abstract in the format of PDF, Word, PowerPoint, jpg, or png, after a manuscript is accepted for publication. For preparing a Professional Graphical Abstract, please click [here](#).



Presentation of the article

Main Format

First page of the manuscripts must be properly identified by the title and the name(s) of the author(s). It should be typed in Times New Roman (font sizes: 17pt in capitalization for the title, 10pt for the section headings in the body of the text and the main text, double spaced, in A4 format with 2cm margins (both doc./docx formats). All pages and lines of the main text should be numbered consecutively throughout the manuscript. Abbreviations in the article title are not allowed. Manuscripts should be arranged in the following order:

1. **TITLE** (brief, attractive and targeted)
2. **Name(s) and Affiliation(s) of author(s)** (including post code and corresponding Email)
3. **ABSTRACT**
4. **Key words** (separate by semicolons; or comma,)
5. **Abbreviations** (those used throughout the manuscript)
6. **INTRODUCTION** (clear statement of the problem, the relevant literature on the subject, and the proposed approach or solution)
7. **MATERIAL AND METHOD** (should be complete enough to allow experiments to be reproduced)
8. **RESULTS**
9. **DISCUSSION**
10. **CONCLUSION**
11. **DECLARATIONS** (Acknowledgements, Consent to publish, Competing interests, Authors' contributions, and Availability of data etc.)
12. **REFERENCES**
13. **Tables**
14. **Figures**
15. **Graphs**

Results and Discussion can be presented jointly.

Discussion and Conclusion can be presented jointly.

Article Sections Format

Title should be a brief phrase describing the contents of the paper. The first letter of each word in title should use upper case. The Title Page should include the author(s)'s full names and affiliations, the name of the corresponding author along with phone and e-mail information. Present address (es) of author(s) should appear as a footnote.

Abstract should be informative and completely self-explanatory, briefly present the topic, state the scope of the experiments, indicate significant data, and point out major findings and conclusions. The abstract should be 150 to 300 words in length. Complete sentences, active verbs, and the third person should be used, and the abstract should be written in the past tense. Standard nomenclature should be used and abbreviations should be avoided. No literature should be cited.

Following the abstract, about 3 to 8 **key words** that will provide indexing references should be listed.

Introduction should provide a clear statement of the problem, the relevant literature on the subject, and the proposed approach or solution. It should be understandable to colleagues from a broad range of scientific disciplines.

Material and Method should be complete enough to allow experiments to be reproduced. However, only truly new procedures should be described in detail; previously published procedures should be cited, and important modifications of published procedures should be mentioned briefly. Capitalize trade names and include the manufacturer's name and address. Subheadings should be used. Methods in general use need not be described in detail. The **ethical approval** for using human and animals in the researches should be indicated in this section with a separated title.

Results should be presented with clarity and precision. The results should be written in the past tense when describing findings in the author(s)'s experiments. Previously published findings should be written in the present tense. Results should be explained, but largely without referring to the literature. In case of the effectiveness of a particular drug or other substances as inhibitor in biological or biochemical processes, the results should be provided as **IC₅₀** (**half maximal inhibitory concentration**) or similar appropriate manner.

Discussion should interpret the findings in view of the results obtained in this and in past studies on this topic. State the conclusions in a few sentences at the end of the paper. The Results and Discussion sections can include subheadings, and when appropriate, both sections can be combined.

Conclusion should be brief and tight about the importance of the work or suggest the potential applications and extensions. This section should not be similar to the Abstract content.

Declarations including Acknowledgements, Author contribution, Competing interests, Consent to publish, and Availability of data etc.

Tables should be kept to a minimum and be designed to be as simple as possible. Tables are to be typed double-spaced throughout, including headings and footnotes. Each table should be on a separate page, numbered consecutively in Arabic numerals and supplied with a heading and a legend. Tables should be self-explanatory without reference to the text. The details of the methods used in the experiments should preferably be described in the legend instead of in the text. The same data should not be presented in both table and graph forms or repeated in the text.

Figure legends should be typed in numerical order on a separate sheet. Graphics should be prepared using applications capable of generating high resolution GIF, TIFF, JPEG or PowerPoint before pasting in the Microsoft Word manuscript file. Use Arabic numerals to designate figures and upper case letters for their parts (Figure 1). Begin each legend with a title and include sufficient description so that the figure is understandable without reading the text of the manuscript. Information given in legends should not be repeated in the text.

Declarations

Please ensure that the sections: Ethics (and consent to participate, if any), Acknowledgements, Author contribution, Competing interests, Consent to publish, Availability of data and materials are included at the end of your manuscript in a Declarations section.

Acknowledgements

We encourage authors to include an Acknowledgements section. Please acknowledge anyone who contributed towards the study by making substantial contributions to conception, design, acquisition of data, or analysis and interpretation of data, or who was involved in drafting the manuscript or revising it critically for important intellectual content, but who does not meet the criteria for authorship. Please also include their source(s) of funding. Please also acknowledge anyone who contributed materials essential for the study. Authors should obtain permission to acknowledge from all those mentioned in the Acknowledgements. Please list the source(s) of funding for the study, for each author, and for the manuscript preparation in the acknowledgements section. Authors must describe the role of the funding body, if any, in study design; in the collection, analysis, and interpretation of data; in the writing of the manuscript; and in the decision to submit the manuscript for publication.

Author contribution

For manuscripts with more than one author, JLSB require an Author Contributions section to be placed after the Acknowledgements section. An 'author' is generally considered to be someone who has made substantive intellectual contributions to a published study. To qualify as an author one should 1) have made substantial contributions to conception and design, or acquisition of data, or analysis and interpretation of data; 2) have been involved in drafting the manuscript or revising it critically for important intellectual content; and 3) have given final approval of the version to be published. Each author should have participated sufficiently in the work to take public responsibility for appropriate portions of the content. Acquisition of funding, collection of data, or general supervision of the research group, alone, does not justify authorship. **We suggest the following format/example** (please use initials to refer to each author's contribution): AB carried out the molecular genetic studies, participated in the sequence alignment and drafted the manuscript. JY carried out the immunoassays. MT participated in the sequence alignment. ES participated in the design of the study and performed the statistical analysis. FG conceived of the study, and participated in its design and coordination and helped to draft the manuscript. All authors read and approved the manuscript. For authors that equally participated in a study please write 'All/Both authors contributed equally to this work.' Contributors who do not meet the criteria for authorship should be listed in an acknowledgements section.

Competing interests

Competing interests that might interfere with the objective presentation of the research findings contained in the manuscript should be declared in a paragraph heading "Competing interests" (after Acknowledgment or Author Contribution sections). Examples of competing interests are ownership of stock in a company, commercial grants, board membership, etc. If there is no competing interest, please use the statement "The authors declare that they have no competing interests.". *Journal of Life Science and Biomedicine* adheres to the definition of authorship set up by [The International Committee of Medical Journal Editors \(ICMJE\)](#). According to the ICMJE authorship criteria should be based on 1) substantial contributions to conception and design of, or acquisition of data or analysis and interpretation of data, 2) drafting the article or revising it critically for important intellectual content and 3) final approval of the version to be published. Authors should meet conditions 1, 2 and 3. It is a requirement that all authors have been accredited as appropriate upon submission of the manuscript. Contributors who do not qualify as authors should be mentioned under Acknowledgements.

Consent to publish

Please include a 'Consent for publication section in your manuscript. If your manuscript contains any individual person's data in any form (including individual details, images or videos), consent to publish must be obtained from that person, or in the case of children, their parent or legal guardian. All presentations of case reports must have consent to publish. You can use your institutional consent form or our consent form if you prefer. You should not send the form to us on submission, but we may request to see a copy at any stage (including after publication). If your manuscript does not contain any individual persons data, please state "Not applicable" in this section.

Change in authorship

We do not allow any change in authorship after provisional acceptance. We cannot allow any addition, deletion or change in sequence of author name. We have this policy to prevent the fraud.

Data deposition

Nucleic acid sequences, protein sequences, and atomic coordinates should be deposited in an appropriate database in time for the accession number to be included in the published article. In computational studies where the sequence information is unacceptable for inclusion in databases because of lack of experimental validation, the sequences must be published as an additional file with the article.

REFERENCES

A JLSB reference style for [EndNote](#) may be found [here](#). However, we prefer [Vancouver](#) referencing style that is often used in medicine and the natural sciences. Uniform requirements for manuscripts submitted to Biomedical Journals, published by International Committee of Medical Journal Editors, includes a list with examples of references https://www.nlm.nih.gov/bsd/uniform_requirements.html in the *Vancouver* style.

References should be numbered consecutively and cited in the text by number in square brackets [1, 2] instead of parentheses (and not by author and date). References should not be formatted as footnotes. Avoid putting personal communications and unpublished observations as references. All the cited papers in the text must be listed in References. All the papers in References must be cited in the text. Where available, URLs for the references should be provided.

Examples (at the text, blue highlighted)

Smit [1] ...; Smit and Janak [2]...; Nurai et al. [3] reported that ; ... [1], --- [2, 3], --- [3-7]. The references at the end of this document are in the preferred referencing style. Give all authors' names; do not use "et al." unless there are six authors or more. Use a space after authors' initials. Papers that have not been published should be cited as "unpublished". Papers that have been accepted for publication, but not yet specified for an issue should be cited as "to be published". Papers that have been submitted for publication should be cited as "submitted for publication". Capitalize only the first word in a paper title, except for proper nouns and element symbols. For papers published in translation journals, please give the English citation first, followed by the original foreign-language citation.

Acceptable Examples (at References section)

For Journals:

1. Hasan V, Sri Widodo M and Semedi B. 2015. Oocyte diameter distribution and fecundity of Javaen Barb (*Systemus Orphoides*) at the start of rainy season in Lenteng River, East Java, Indonesia insurance. J. Life Sci Biomed, 5(2): 39-42.
2. Karen KS, Otto CM. 2007. Pregnancy in women with valvular heart disease. Heart. 2007 May; 93(5): 552-558.
3. Doll MA, Salazar-González RA, Bodduluri S, Hein DW. Arylamine N-acetyltransferase 2 genotype-dependent N-acetylation of isoniazid in cryopreserved human hepatocytes. Acta Pharm Sin B, 2017; 7(4):517-522.

For In press manuscripts (maximum 2):

Hasan V, Sri Widodo M and Semedi B. 2015. Oocyte Diameter Distribution and Fecundity of Javaen Barb (*Systemus Orphoides*) at the Start of Rainy Season in Lenteng River, East Java, Indonesia insurance. In press.

For symposia reports and abstracts:

Cruz EM, Almatar S, Aludul EK and Al-Yaqout A. 2000. Preliminary Studies on the Performance and Feeding Behaviour of Silver Pomfret (*Pampus argentens euphrasen*) Fingerlings fed with Commercial Feed and Reared in Fibreglass Tanks. Asian Fisheries Society Manila, Philippine 13: 191-199.

For Conference:

Skinner J, Fleener B and Rinchiuso M. 2003. Examining the Relationship between Supervisors and Subordinate Feeling of Empowerment with LMX as A Possible Moderator. 24th Annual Conference for Industrial Organizational Behavior.

For Book:

Russell, Findlay E, 1983. Snake Venom Poisoning, 163, Great Neck, NY: Scholium International. ISBN 0-87936-015-1.

For Web Site:

Bhatti SA and Firkins JT. 2008. http://www.ohioline.osu.edu/sc1156_27.html.

Nomenclature and Abbreviations

Nomenclature should follow that given in NCBI web page and Chemical Abstracts. Standard abbreviations are preferable. If a new abbreviation is used, it should be defined at its first usage. Abbreviations should be presented in one paragraph, in the format: "term: definition". Please separate the items by ";".

E.g. ANN: artificial neural network; CFS: closed form solution; ...

Abbreviations of units should conform with those shown below:

Decilitre	dl	Kilogram	kg
Milligram	mg	hours	h
Micrometer	mm	Minutes	min
Molar	mol/L	Mililitre	ml
Percent	%	.	

Other abbreviations and symbols should follow the recommendations on units, symbols and abbreviations: in "A guide for Biological and Medical Editors and Authors (the Royal Society of Medicine London 1977). Papers that have not been published should be cited as "unpublished". Papers that have been accepted for publication, but not yet specified for an issue should be cited as "to be published". Papers that have been submitted for publication should be cited as "submitted for publication".

Formulae, numbers and symbols

1. Typewritten formulae are preferred. Subscripts and superscripts are important. Check disparities between zero (0) and the letter O, and between one (1) and the letter I.
2. Describe all symbols immediately after the equation in which they are first used.
3. For simple fractions, use the solidus (/), e.g. 10 /38.
4. Equations should be presented into parentheses on the right-hand side, in tandem.
5. Levels of statistical significance which can be used without further explanations are *P < 0.05, **P < 0.01, and ***P < 0.001.
6. In the English articles, a decimal point should be used instead of a decimal comma.
7. Use Symbol fonts for "±"; "≤" and "≥" (avoid underline).
8. In chemical formulae, valence of ions should be given, e.g. Ca²⁺ and CO₃²⁻, not as Ca⁺⁺ or CO₃.
9. Numbers up to 10 should be written in the text by words. Numbers above 1000 are recommended to be given as 10 powered x.
10. Greek letters should be explained in the margins with their names as follows: Αα - alpha, Ββ - beta, Γγ - gamma, Δδ - delta, Εε - epsilon, Ζζ - zeta, Ηη - eta, Θθ - theta, Ιι - iota, Κκ - kappa, Λλ - lambda, Μμ - mu, Νν - nu, Ξξ - xi, Οο - omicron, Ππ - pi, Ρρ - rho, Σσ - sigma, Ττ - tau, Υυ - ipsilon, Φφ - phi, Χχ - chi, Ψψ - psi, Ωω - omega. Please avoid using math equations in Word whenever possible, as they have to be replaced by images in xml full text.

Review/Decisions/Processing/Policy

Firstly, all manuscripts will be checked by [Docol@c](#), a plagiarism finding tool. The received papers with plagiarism rate of more than 30% will be rejected. Manuscripts that are judged to be of insufficient quality or unlikely to be competitive enough for publication will be returned to the authors at the initial stage. The remaining manuscripts go through a single-blind review process by external reviewers selected by section editor of JLSB, who are research workers specializing in the relevant field of study. One unfavourable review means that the paper will not be published and possible decisions are: accept as is, minor revision, major revision, or reject. The corresponding authors should submit back their revisions within 14 days in the case of minor revision, or 30 days in the case of major revision. Manuscripts with significant results are typically published at the highest priority. The editor who received the final revisions from the corresponding authors shall not be hold responsible for any mistakes shown in the final publication.

The submissions will be processed free of charge for invited authors, authors of hot papers, and corresponding authors who are editorial board members of the *Journal of Life Science and Biomedicine*. This journal encourage the academic institutions in low-income countries to publish high quality scientific results, free of charges.

Plagiarism

Manuscripts are screened for plagiarism by [Docol@c](#), before or during publication, and if found (more than 30% duplication limit) they will be rejected at any stage of processing. If we discovered accidental duplicates of published article(s) that are determined to violate our journal publishing ethics guidelines (such as multiple submission, bogus claims of authorship, plagiarism, fraudulent use of data or the like), the article will be "Withdrawn" from SCIENCELINE database. Withdrawn means that the article content (HTML and PDF) is removed and replaced with a HTML page and PDF simply stating that the article has been withdrawn according to the [Scienceline Policy](#) on Published Article Withdrawal.

Date of issue

All accepted articles are published bimonthly around 25th of January, March, May, July, September and November, each year in full text on the internet.

The OA policy

Journal of Life Science and Biomedicine is an open access journal which means that all content is freely available without charge to the user or his/her institution. Users are allowed to read, download, copy, distribute, print, search, or link to the full texts of the articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author. This is in accordance with the [BOAI definition of Open Access](#).

Submission Preparation Checklist

- Authors are required to check off their submission's compliance with all of the following items, and submissions may be returned to authors that do not adhere to the following guidelines.
- The submission has not been previously published, nor is it before another journal for consideration (or an explanation has been provided in Comments to the Editor).
- The submission file is in Microsoft Word, RTF, or PDF document file format. Where available, URLs for the references have been provided.
- The text is single-spaced; uses a 12-point font; and all illustrations, figures, and tables are placed within the text at the appropriate points, rather than at the end. The text adheres to the stylistic and bibliographic requirements outlined in the Author Guidelines.

Paper Submission Flow



SCIENCELINE PUBLISHING CORPORATION


Scienceline Publication Ltd is a limited liability non-profit non-stock corporation incorporated in Turkey, and also is registered in Iran. Scienceline journals that concurrently belong to many societies, universities and research institutes, publishes internationally peer-reviewed open access articles and believe in sharing of new scientific knowledge and vital research in the fields of life and natural sciences, animal sciences, engineering, art, linguistic, management, social and economic sciences all over the world. Scienceline journals include:

Online Journal of Animal and Feed Research



ISSN 2228-7701; Bi-monthly
[View Journal](#) | [Editorial Board](#)
 Email: editors@ojaf.ir
[Submit Online >>](#)

Journal of Civil Engineering and Urbanism




ISSN 2252-0430; Bi-monthly
[View Journal](#) | [Editorial Board](#)
 Email: ojceu@ojceu.ir
[Submit Online >>](#)

Journal of Life Sciences and Biomedicine




ISSN: 2251-9939; Bi-monthly
[View Journal](#) | [Editorial Board](#)
 Email: editors@jlsb.science-line.com
[Submit Online >>](#)

Asian Journal of Medical and Pharmaceutical Researches



ISSN: 2322-4789; Quarterly
[View Journal](#) | [Editorial Board](#)
 Email: editor@ajmpr.science-line.com
[Submit Online >>](#)

Journal of World's Poultry Research



ISSN: 2322-455X; Quarterly
[View Journal](#) | [Editorial Board](#)
 Email: editor@jwpr.science-line.com
[Submit Online >>](#)

World's Veterinary Journal




ISSN: 2322-4568; Quarterly
[View Journal](#) | [Editorial Board](#)
 Email: editor@wjv.science-line.com
[Submit Online >>](#)

Journal of Educational and Management Studies




ISSN: 2322-4770; Quarterly
[View Journal](#) | [Editorial Board](#)
 Email: info@jems.science-line.com
[Submit Online >>](#)

Journal of World's Electrical Engineering and Technology



ISSN: 2322-5114; Irregular
[View Journal](#) | [Editorial Board](#)
 Email: editor@jweet.science-line.com
[Submit Online >>](#)

Journal of Art and Architecture Studies



ISSN: 2383-1553; Irregular
[View Journal](#) | [Editorial Board](#)
 Email: jaas@science-line.com
[Submit Online >>](#)

Asian Journal of Social and Economic Sciences



ISSN: 2383-0948; Quarterly
[View Journal](#) | [Editorial Board](#)
 Email: ajses@science-line.com
[Submit Online >>](#)

Journal of Applied Business and Finance Researches



ISSN: 2382-9907; Quarterly
[View Journal](#) | [Editorial Board](#)
 Email: jabfr@science-line.com
[Submit Online >>](#)

Scientific Journal of Mechanical and Industrial Engineering



ISSN: 2383-0980; Quarterly
[View Journal](#) | [Editorial Board](#)
 Email: sjmie@science-line.com
[Submit Online >>](#)