

A Review on Challenges and Opportunities of Water Hyacinth

Tadlo Awoke Mengesha✉✉

Department of Animal Production and Extension, College of Veterinary Medicine and Animal Sciences, University of Gondar, P.O.Box 196 Gondar, Ethiopia

✉Corresponding Email: mebratuaw@gmail.com

ABSTRACT

This paper reviews the major existing challenges and their possible mitigation of water hyacinth. Water hyacinth (*Eichhornia crassipes*) is a cosmopolitan invasive aquatic plant which can tolerate a wide range of environmental conditions such as temperature, humidity, illumination, pH, salinity, wind, current and drought. The plant is morphologically plastic with a rapid mode of vegetative propagation that makes it well adapted to a long distance of dispersal and colonization under diverse ecological conditions. Origin of water hyacinth on the African continent was first reported in Egypt between 1879 and 1893; in South Africa in 1908, Zimbabwe in 1937, Zaire and Sudan in 1957, Senegal in 1964, Ethiopia in 1965, Nigeria in 1983 and Uganda in 1987. Water hyacinth has a multitude of direct and indirect effects. The main problems arising from the growth of Water Hyacinth in thick mats are: an enormous water loss through evapotranspiration, that alters the water balance of entire regions; the impediment to water flow, that increases sedimentation, causing flooding and soil erosion; the obstruction of navigation; hampering fishing and dramatically reducing the catch and the source of food and income for local populations; a drastic change in the physical and chemical properties of water and in the environment in the water bodies invaded, with detrimental effects on plants and animals; the reduction of the activity of electrical power stations, jeopardizing the power supply of the country and a serious threaten to agricultural production, following the blockage of irrigation canals and drainage systems. Although water hyacinth is seen in many countries including Ethiopia as a weed and is responsible for many of the problems outlined earlier in this fact sheet, many individuals, groups and institutions have been able to turn the problem around and find useful applications for the plant. Water hyacinth has received much attention in recent years due to its potential benefits as animal fodder, aqua feed, water purification, fertilizer, biogas production. There are several popular control mechanisms for preventing the spread of, or eradication of water hyacinth. The three main mechanisms used such as biological, chemical and physical control.

Review

PII: S225199391700009-7

Rec. 05 Aug. 2017
Acc. 01 Sep. 2017
Pub. 25 Sep. 2017

Keywords

Aquatic weed,
Challenges,
Ecology,
Management,
Water hyacinth

INTRODUCTION

Water Hyacinth belongs to Kingdom-Plantae, Order-Commelinids, Family-Pontederiaceae, Genus- Eichhornia and Species-crassipes. Water hyacinth (*Eichhornia crassipes*) is a cosmopolitan invasive aquatic plant which can tolerate a wide range of environmental conditions such as temperature, humidity, illumination, pH, salinity, wind, current and drought. The plant is morphologically plastic with a rapid mode of vegetative propagation that makes it well adapted to a long distance of dispersal and colonization under diverse ecological conditions. It is one of the most prolific aquatic plant which spreads at an alarming rate. It has spikes of light blue flowers and green color roundish leaves with inflated bladder like petioles. Water hyacinth (*Eichhornia crassipes*), native to South America, but now an environmental and social menace throughout the old world tropics, affects the environment and humans in

diverse ways. Most of these are detrimental, although some are beneficial or potentially useful. Many of these effects are due to its potential to grow rapidly and produce enormous amounts of biomass, thereby covering extensive areas of naturally open water [1]. The extremely rapid rate of proliferation of *Eichhornia crassipes* results reduced penetration of dissolved oxygen in water body, change in water chemistry, disruption of aquatic flora and increased rate of water loss due to evapotranspiration. *Eichhornia crassipes* is very popular recently as animal feed, aqua feed, water purification, fertilizer, biogas production, even food for human and other products [2, 3]. Therefore, it is considered as a serious threat to biodiversity and recently massive attention has been given to its harvesting for use as alternative plant protein source for livestock.

Origin and distribution of water hyacinth

Water hyacinth (*Eichhornia crassipes*) species is common and widely distributed all over the world. Water hyacinth are a free-floating perennial aquatic plant native to tropical and sub-tropical South America; with bright green, waxy leaves and attractive, violet flowers that have yellow stripes on the banner petals. It is widely reported that water hyacinth is indigenous to Brazil having first been described from wild plants collected from Francisco River in 1824. By the early 1990s it had spread to virtually every country in the continent [4]. It is believed that the water hyacinth was introduced first into the United States at the World's Industrial and Cotton Centennial Exposition of 1884-1885 in Louisiana. Similarly Water Hyacinth (*Eichhornia crassipes* (Martius) Solms-Laubach), which is native to the Amazon basin, Brazil [5, 6] became widespread throughout the world, also due to its attractive appearance. It is commercially available as an ornamental for ponds. Its spread started with its deliberate introduction into North America from Brazil, in the late nineteenth century, as an ornamental in ponds and subsequently escaped cultivation [5][6]. At present it occurs as a weed throughout tropical and subtropical regions of the world, including North and South America, Africa, Asia, Australia and New Zealand.

Water hyacinth has invaded freshwater systems in over 50 countries on five continents; it is especially pervasive throughout Southeast Asia, the southeastern United States, central and western Africa, and Central America [7-10]. On the African continent, water hyacinth was first reported in Egypt between 1879 and 1893; in South Africa in 1908; Zimbabwe (1937); Zaire and Sudan (1957); Senegal (1964); Ethiopia (1965); Nigeria (1983) and Uganda (1987). Penfound and Earle [4] reported that Water hyacinth spread through fragmentation of established plants and may re-sprout from rhizomes or germinate from seeds. Dispersal also occurs by water-borne seeds and by seeds that stick to the feet of birds. Migratory birds may be important in long distance dispersal [11]. The major means of dispersal, and the most difficult to control, is active transport by people who, ignorant of its impacts, seek to propagate it in other ponds and lakes. Humans also contribute to its spread in some areas by using the plant as a packing material and as cushions in boats [11].

Although exotic aquatic weeds have been reported to be present in Africa since the end of the nineteenth century [12] they started infesting massively African freshwater bodies during the early 1950s [13] and rapidly spread in many countries. The growth of these weeds is extremely fast and this allows them to develop huge infestations in areas where they had not been reported only a few years earlier. This plants invaded lakes, ponds, rivers, canals and agricultural fields, becoming noxious weeds. The damage to the environment and the economy is enormous, having a disrupting impact on agriculture, fisheries, production of electricity, transportation, health, means of sustenance, living conditions and social structure. Water Hyacinth, Water Lettuce and Water Fern pose the most serious problems.

Botanists and gardeners carry plants with them in their travels, and experts suspect that this is how the water hyacinth came to East Africa in the 1980s. Due to its attractive flowers; it was probably brought over as an ornamental for garden ponds [14]. The consensus is that Water Hyacinth entered Lake Victoria from Rwanda via the river Kagera [15]. The exact time and place of introduction has been debated, but the plant is native to South America, and therefore reached Lake Victoria due to human activity. It has spread prolifically, due to lack of natural enemies, an abundance of space, agreeable temperature conditions, and abundant nutrients [16]. It increased rapidly between 1992 -1998, was greatly reduced by 2001, and has since resurged to a lesser degree. Management techniques include (hyacinth-eating) insect controls and manual beach cleanup efforts [17].

Water hyacinth, *Eichhornia crassipes*, was first reported in Ethiopia in 1965 in Koka Lake and the Awash River. The Koka Lake (also known as Lake Gelila) is a reservoir in south-central part of Ethiopia created by the construction of the Koka Dam across the Awash River [18]. Since the first invasion, in the above reservoir, the weed has also been found in Lake Ellen and other rivers but never in the upper catchment of the Blue Nile until most recently. Sadly, in September 2011, it was officially recognized that one of the top ten ecologically dangerous and worst invasive weed, water hyacinth (*Eichhornia crassipes*), infested Lake Tana [19]. Except speculation, the exact source of water hyacinth

infestation of Lake Tana is not known. The first infestations were found near the mouth of the Megech River on the northern shores of the lake. This noxious weed is now abundant along 40 km of the northern and north-eastern shoreline of the Lake, posing a significant threat to livelihoods, biodiversity, tourism and the general ecological health of the Lake Tana ecosystem. The infestation is greater in the northern tip of the Lake at Demebia district and is minimized towards the eastern tip of Libo-Kemkem district. A survey by a team of experts from the BoEPLAU conducted in September 2012 puts the infestation level at 20,000 hectares compared to 4,000 hectares in 2011. The 2014 survey estimate of water hyacinth coverage was nearly 50 000 ha shore area of the lake, greater than doubled the 2012 coverage. Anteneh [20] Survey report revealed that water Hyacinth in Lake Tana the most devastating area coverage by the weed was observed at Megech River mouth extended both east and north direction with estimated ca. 34 500 ha (3162 ha thick, 2591 ha intermediate and 28687 ha scattered) and widely distribution of daughter plants observed that moved forward by the assistance of the wave.

DISCUSSION

Morphology and biology of water hyacinth

Water hyacinth is a perennial herbaceous plant is a floating freshwater hydrophyte. It belongs to the Family Pontederiaceae and all the species in the Genus Eichhornia are aquatic. Water hyacinth (*Eichhornia crassipes*) tends to form mats on the water surface. Sometimes water hyacinth can be found growing in muddy soils near the edge of an aquatic system. The leaves are arranged in a rosette. The plant is morphologically very plastic with a rapid mode of vegetative propagation which makes it well adapted to long distance dispersal and successful colonization of diverse ecological niches. It is one of the most prolific aquatic plants which spread at an alarming rate having spikes of large blue flowers and roundish leaves with inflated bladder - like petioles. The leaf stem usually is somewhat to completely swollen and filled with spongy tissue and thus acts as afloat. The blade of the leaf is oval to round and usually much smaller than the leaf stem. The common water hyacinth (*Eichhornia crassipes*) is vigorous growers known to double their population in two weeks. Water hyacinth grows rapidly. Growth of more than one tone of dry matter per day per hectare is not uncommon. One plant may be able to produce enough growth to cover 600 square meters in one year. Infestation breaks up in to "rafts" that drift wherever the winds and currents take them, rapidly infesting entire river systems [11].

Water Hyacinth shows considerable variation in both leaf and flower form and colour, also depending on the age of the plant. The flowers are bluish purple, large and self-fertile. The seeds are produced in large numbers and are contained in capsules, each capsule containing up to 300 seeds [21]. The seeds can remain viable for 5-20 years [22]. The plant can also reproduce vegetatively through the production of horizontal stolns.

Negative consequences of water hyacinth

Water hyacinth has a multitude of direct and indirect effects on almost all aspects of human life once a water body on which man so much depends is invaded and covered by the weed mats [23]: fisheries; water supply; hydroelectric power generation; human health; agriculture; transport; biodiversity; evapotranspiration and increased cost of water treatment are some of the adverse effects. There are already negative environmental and economic impacts caused by water hyacinth on Lake Tana and these include impairing and blocking of fishing grounds and fish breeding areas, reduced fish catches and fish quality, reduced tourism potential, reduced recession agricultural production and blockage of irrigation canals, loss of pasture lands and blocking of fishers movements could also be mentioned. Health impacts include increase in vector-borne diseases such as malaria, bilharzias and health related hazards from leeches and reptiles. Environmental impacts include deterioration in water quality, water loss through evapo-transpiration and a decline of aquatic biodiversity. Water hyacinth can cause a variety of problems when its rapid mat-like proliferation covers areas of fresh water. Some of the common problems are listed below:

Fish production. Once the water body is covered by the water hyacinth fishing activities will be curtailed as landing sites would be inaccessible. Furthermore breeding sites will be reduced and fishermen take longer to reach fishing grounds. According Kateregga and Sterner [24] report stated that water hyacinth mats invaded fishing grounds and blocked waterways. For the individual fisherman, the hyacinth mats reduced their catch by covering fishing grounds, delaying access to markets due to loss of output, increasing fishing costs due to the time and effort spent clearing waterways, forcing translocation, and causing loss of nets. Mailu's [25] report cited the declines of fish production by 14 percent, 37 percent, and 59 percent in the catches of *Oriochromis* (a large genus of tilapia), *Clarias*

(a genus of catfish), and *Mormyrus* (a genus of bottom-feeding breams), respectively, in the Kenyan section of the Lake Victoria. According to Twongo [11] noted that the weed mats sealed off breeding, nursery, feeding, and fishing grounds for various inshore fish species, like tilapia and young Nile perch. The mats also had detrimental effects by blocking light, severely reducing oxygen levels, and allowing poisonous gases, such as ammonia and hydrogen sulfide, to accumulate.

Kateregga and Sterner [24] stated that the effect of the water hyacinth on the catch ability of fish in the Kenya, Tanzania, and Uganda fisheries of Lake Victoria by incorporating the water hyacinth biomass as a negative factor in the catch ability coefficient. The results indicate that the catch ability of fish in the Lake Victoria fisheries was reduced by a factor of 2–45 percent during the period when the lake was highly infested by the water hyacinth. The relationship between water hyacinth infestation and fish production is inversely. Kateregga and Sterner [24] confirmed that the larger reduction in the catch ability of fish in Kenya's section was explained by the high abundance of water hyacinth mats in this area, compared, to Tanzania and Uganda.

Weed problem. The fast growth of Water Hyacinth allowed the plant to build huge populations in its ranges of introduction, developing dense mats on the surface of the water and becoming a major weed problem. It is considered the worst aquatic weed in the world. The rapid increase and spread of the plant into new areas is due particularly to its vegetative reproduction, a single plant being able to develop very rapidly a significant infestation. Moving easily with water currents, winds or other accidental means, such as fishing nets and boats, the plant invaded rivers, canals, ponds, lakes, dams and other freshwater bodies. Water hyacinth acts as a weed in paddy rice by interfering with germination and establishment [1]. The main problems arising from the growth of Water Hyacinth in thick mats are (a) an enormous water loss through evapotranspiration, that alters the water balance of entire regions; (b) the impediment to water flow, that increases sedimentation, causing flooding and soil erosion; (c) the obstruction of navigation; (d) hampering fishing and dramatically reducing the catch and the source of food and income for local populations; (e) a drastic change in the physical and chemical properties of water and in the environment in the water bodies invaded, with detrimental effects on plants and animals; (f) the reduction of the activity of electrical power stations, jeopardizing the power supply of the country; and (g) a serious threaten to agricultural production, following the blockage of irrigation canals and drainage systems. The economy of the countries concerned was therefore seriously affected in many aspects. This weed represents an environmental problem as well and indirectly a public health problem, since it may create a microhabitat suitable for the breeding of many vectors of human diseases and for hosting poisonous snakes.

Water Hyacinth is the principal aquatic weed in Africa. It is a weed problem in many countries, especially in Egypt and East, West and South Africa. This plant appeared in Lake Kyoga in Uganda in the early 1980s and it also occurs in Lake Kwanza and down the Kyoga Nile. In 1988 it was observed in Lake Naivasha in Kenya. However, the most disturbing development in the region during the 1990s was in Lake Victoria. In 1990 dense mats of this weed were found interfering with fishing on the Tanzanian shore and during the same year mats were recorded on the Ugandan shore and around the Sese Islands. In Zimbabwe the plant attained the status of noxious weed in the 1980s. It was first recorded in Lake Kariba in 1994 and by 1996 it infested over 200 ha. In the early 1990s Water Hyacinth was also found in the Pangani River in Tanzania and the lower reaches of the Kagera River in Rwanda. In Ethiopia around in September 2011, it was officially recognized that one of the top ten ecologically dangerous and worst invasive weed, water hyacinth (*Eichhornia crassipes*), infested Lake Tana [19]. This plants invaded lakes, ponds, rivers, canals and agricultural fields, becoming noxious weeds.

Ecological impact. That is just one of its ecological impacts. Water hyacinth also reduces biological diversity, impacts native submersed plants, alters immersed plant communities by pushing away and crushing them, and also alter animal communities by blocking access to the water and/or eliminating plants the animals depend on for shelter and nesting. In Lake Victoria, African fishermen have noted that, in areas where there is much water hyacinth infestation, the water is still and warm and the fish disappear [26]. They also complain that crocodiles and snakes have become more prevalent.

The physical problems brought about by water hyacinth are now common knowledge. Water hyacinth mats clog waterways, making boating, fishing and almost all other water activities impossible. Many large hydropower schemes are suffering from the effects of water hyacinth.

According to Makhanu [27] revealed that Water supply will be affected as intake works would be clogged and the irrigation canals will be clogged or their hydraulic efficiency drastically reduced. Transport by ships or boats will be hindered. Also, evapotranspiration is increased as loss of biodiversity in the water body covered by the water

hyacinth. The cost of purifying water tainted by water hyacinth will be increased tremendously. Hydroelectric power production will be affected since turbines would be clogged resulting into expensive repair, overhaul and maintenance. Human health will be affected in many ways: shoreline mats are habitats for certain snails (schistosomia vectors) and mosquitoes which spread malaria. Agriculture will be adversely affected [27].

Hindrance to water transport. Access to harbors and docking areas can be seriously hindered by mats of water hyacinth. Canals and freshwater rivers can become impassable as they clog up with densely intertwined carpets of the weed. It is also becoming a serious hazard to lake transport on Lake Victoria as large floating islands of water hyacinth form, while many of the inland waterways of south East Asia have been all but abandoned. Water hyacinth mats are difficult or impossible to penetrate with boats, and even small mats regularly foul boat propellers. This can have a severe effect on transport, especially where water transport is the norm [1].

Blockage of irrigation, hydropower and water supply systems. Many large hydropower schemes are suffering from the effects of water hyacinth. The Owen Falls hydropower scheme at Jinja on Lake Victoria is a victim of the weeds rapid reproduction rates and an increasing amount of time and money has to be invested in clearing the weed to prevent it entering the turbine and causing damage and power interruptions. Water hyacinth is now a major problem in some of the world's major dams - the Kariba dam which straddles the Zambia-Zimbabwe border on the Zambezi River and feeds Harare has pronounced infestations of the weed. Cock [1] revealed that Water hyacinth replaces existing aquatic plants, and develops floating mats of interlocked water hyacinth plants, which are colonized by several semi-aquatic plant species. As succession continues, floating mats dominated by large grasses may drift away or be grounded. This process can lead to rapid and profound changes in wetland ecology, e.g. shallow areas of water will be converted to swamps. In slow-moving water bodies, water hyacinth mats physically slow the flow of water, causing suspended particles to be precipitated, leading to silting. The reduced water flow can also cause flooding and adversely affect irrigation schemes. Water hyacinth acts as a weed in paddy rice by interfering with germination and establishment.

Reduction of biodiversity. Where water hyacinth is prolific, other aquatic plants have difficulty in surviving. This causes an imbalance in the aquatic micro-ecosystem and often means that a range of fauna that relies on a diversity of plant life for its existence will become extinct. Diversity of fish stocks is often affected with some benefiting and others suffering from the proliferation of water hyacinth. People often complain of localized water quality deterioration. This is of considerable concern where people come to collect water and to wash. Cock [1] stated that Water hyacinth has direct effects upon water chemistry. It can absorb large amounts of nitrogen and phosphorus, other nutrients and elements. It is this ability to pick up heavy metals which has led to the suggestion that water hyacinth could be used to help clean industrial effluent in water. By absorbing and using nutrients, water hyacinth deprives phytoplankton of them. This leads to reduced phytoplankton, zooplankton and fish stocks. Conversely, as the large amounts of organic material produced from senescent water hyacinth decompose, this leads to oxygen deficiency and anaerobic conditions under the floating water hyacinth mats. These anaerobic conditions have been the direct cause of fish death, and changes in the fish community by eliminating most species at the expense of air breathing species.

Increased evapotranspiration. Various studies have been carried out to certain the relationship between aquatic plants and the rate of evapotranspiration compared with evaporation from an open-surfaced water body. Saelthun [28] suggests that the rate of water loss due to evapotranspiration can be as much as 1.8 times that of evaporation from the same surface but free of plants. This has great implications where water is already scarce. It is estimated that the flow of water in the Nile could be reduced by up to one tenth due to increased losses in Lake Victoria from water hyacinth. Water hyacinth is reported to cause substantially increased loss of water by evapotranspiration compared to open water, although this has recently been challenged. Displacement of water by water hyacinth can mean that the effective capacity of water reservoirs is reduced by up to 400 m³ of water per hectare, causing water levels in reservoirs to fall more rapidly in dry periods. Water displacement, siltation of reservoirs and physical fouling of water intakes can have a major impact on hydroelectric schemes [1].

Problems related to fishing. Water hyacinth can present many problems for the fisherman. Access to sites becomes difficult when weed infestation is present, loss of fishing equipment often results when nets or lines

become tangled in the root systems of the weed and the result of these problems is more often than not a reduction in catch and subsequent loss of livelihood. Infestations make access to fishing grounds increasingly time consuming or impossible, while physical interference with nets makes fishing more difficult or impractical. Some fishing communities in West Africa have been abandoned as a direct result of the arrival of water hyacinth [1]. In areas where fishermen eke a meager living from their trade, this can present serious socio-economic problems. Edward [30] stated water hyacinth infestation in Lake Tana is a threat to the fishing industry that supports the livelihood of an estimated 20,000 riparian communities and who are dependent on this natural resource. A reduction in water levels during the dry season (October to May but peak between July and September) caused large amounts of the water hyacinth to die and decomposition of the water hyacinth caused massive algal blooms which affected the taste of the fish with the result that fisher folk were unable to sell their catch. The impacts of this weed are similar to those on Lake Victoria. Fishermen on Lake Victoria have also noted that, in areas where there is much water hyacinth infestation, the water is 'still and warm and the fish disappear. They also complain that crocodiles and snakes. Conversely, as the large amounts of organic material produced from senescent water hyacinth decompose, this leads to oxygen deficiency and anaerobic conditions under the floating water hyacinth mats. These anaerobic conditions have been the direct cause of fish death, and changes in the fish community by eliminating most species at the expense of air breathing species. Stationary mats of water hyacinth also shade out bottom growing vegetation, thereby depriving some species of fish, of food and spawning grounds. The potential impact on fish diversity is enormous.

Micro-habitat for a variety of disease vectors. The diseases associated with the presence of aquatic weeds in tropical developing countries are among those that cause the major public health problems: malaria, schistosomiasis and lymphatic filariasis. Some species of mosquito larvae thrive on the environment created by the presence of aquatic weeds, while the link between schistosomiasis (bilharzia) and aquatic weed presence is well known. Although the statistical link is not well defined between the presence of aquatic weeds and malaria and schistosomiasis, it can be shown that the brughian type of filariasis (which is responsible for a minor share of lymphatic filariasis in South Asia) is entirely linked to the presence of aquatic weeds. Cock [1] reported that water hyacinth encourage the vectors of several human diseases, including the intermediate snail hosts of bilharzia (schistosomiasis) and most mosquito vectors, including those responsible for the transmission of malaria, encephalitis and filariasis. In parts of Africa, water hyacinth mats are reported to provide cover for lurking crocodiles and snakes.

Positive consequence of water hyacinth

Although water hyacinth is seen in many countries including Ethiopia as a weed and is responsible for many of the problems outlined earlier in this fact sheet, many individuals, groups and institutions have been able to turn the problem around and find useful applications for the plant. A water hyacinth composition have more than 95% water, has a fibrous tissue and a high energy and protein content, and can be used for a variety of useful applications. Water hyacinth harvests have been put into valuable uses in several countries. Methods of converting the plant material into valuable products have emerged. Apart from its ornamental value, the plant has been found useful as a source of animal feed [30] as a source of fertilizers for use in agriculture [31, 32], a source of bio- mass energy, a source of raw materials for building, handcraft making, paper and boards. In addition the plant has been found to be useful as a filter worth of solving man created problems of pollution in water bodies. However all the potential uses of the water hyacinth do not promote utilization of the weed to the level that qualifies it as a viable control option [33].

Animal fodder. Studies have shown that the nutrients in water hyacinth are available to ruminants. In Southeast Asia some no ruminant animals are fed rations containing water hyacinth. In China pig farmers boil chopped water hyacinth with vegetable waste, rice bran, copra cake and salt to make a suitable feed. In Malaysia fresh water hyacinth is cooked with rice bran and fishmeal and mixed with copra meal as feed for pigs, ducks and pond fish. Similar practices are much used in Indonesia, the Philippines and Thailand [35]. The high water and mineral content mean that it is not suited to all animals. In Bangladesh, huge amount of *Eichhornia crassipes* are produced due to large number of rivers, ponds, lakes and other water reservoirs. In many coastal areas of the country, *Eichhornia crassipes* is commonly used as forage for cattle either as basal feed resource or supplement to a diet consists of sugarcane, molasses and cereal straws.

The use of water hyacinth for animal feed in developing countries could help solve some of the nutritional problems that exist in these countries. Animal feed is often in short supply and although humans cannot eat water hyacinth directly, they can feed it to cattle and other animals which can convert the nutrient into useful food products for human consumption.

Water hyacinth (*Eichhornia crassipes*) an abundantly available aquatic plant of many country considered menace and the age old tradition believe that if this plan is fed to animal it will affect the health of the animals with severe diarrhea and suffer from salt imbalances. Many authors stated the utilization of Water Hyacinth (*Eichhornia Crassipes*) for livestock feed in different form. Many authors stated the utilization of Water Hyacinth (*Eichhornia Crassipes*) for livestock feed in different form. According to Villadolid and Bunag [35] have given a short but useful review of the uses of water hyacinth for animal, fish and human food. Sharma [36] has also discussed the uses of water hyacinth for livestock food. There is also reference is made to the harvesting of water hyacinth in Bangladesh, India, Indonesia and the Sudan to feed livestock.

Basket work. In the Philippines water hyacinth is dried and used to make baskets and matting for domestic use. The key to a good product is to ensure that the stalks are properly dried before being used. If the stalks still contain moisture then this can cause the product to rot quite quickly. In India, water hyacinth is also used to produce similar goods for the tourist industry. Traditional basket making and weaving skills are used.

Biogas production. The possibility of converting water hyacinth to biogas has been an area of major interest for many years. Conversion of other organic matter, usually animal or human waste, is a well-established small and medium scale technology in a number of developing countries, notably in China and India. The process is one of anaerobic digestion which takes place in a reactor or digester (an air tight container usually sited below ground) and the usable product is methane gas which can be used as a fuel for cooking, lighting or for powering an engine to provide shaft power. The residue from the digestion process provides a fertilizer rich in nutrients.

Water purification. Water hyacinth can be used to aid the process of water purification either for drinking water or for liquid effluent from sewage systems. A drinking water treatment water hyacinth has been used as part of the pretreatment purification step. Clean, healthy plants have been incorporated into water clarifiers and help with the removal of small flock that remain after initial coagulation and flock removal or settling [37]. Water hyacinth roots naturally absorb pollutants, including such toxic chemicals as lead, mercury, and strontium 90 (as well as some organic compounds). The result is a significant decrease in turbidity due to the removal of flock and also slight reduction in organic matter in the water. Water hyacinth is already being used to clean up waste water in small scale sewage treatment plants. Phyto remediation used for removing heavy metals and other pollutants is a newly developed environmental protection technique. In sewage systems, the root structures of water hyacinth (and other aquatic plants) provide a suitable environment for aerobic bacteria to function. Aerobic bacteria feed on nutrients and produce inorganic compounds which in turn provide food for the plants. The plants grow quickly and can be harvested to provide rich and valuable compost. Water hyacinth has also been used for the removal or reduction of nutrients, heavy metals, organic compounds and pathogens from water [30].

Fertilizers. Water hyacinth can be used on the land either as a green manure or as compost. As a green manure it can be either ploughed into the ground or used as mulch. The plant is ideal for composting. After removing the plant from the water it can be left to dry for a few days before being mixed with ash, soil and some animal manure. Microbial decomposition breaks down the fats, lipids, proteins, sugars and starches. The mixture can be left in piles to compost, the warmer climate of tropical countries accelerating the process and producing rich pathogen free compost which can be applied directly to the soil. The compost increases soil fertility and crop yield and generally improves the quality of the soil.

Compost can be made on a large or small scale and is well suited to labour intensive, low capital production. In developing countries where mineral fertilizer is expensive, it is an elegant solution to the problem of water hyacinth proliferation and also poor soil quality. In Sri Lanka water hyacinth is mixed with organic municipal waste, ash and soil, composted and sold to local farmers and market gardeners.

Fish feed. At the same time, the water hyacinth is believed to have promoted fish diversity, particularly smaller species and the young. Mechanisms for this include providing shelter from predators as well as reducing fishing pressure. It enhanced the abundance of lungfish and Haplochromines (riverine "haps") and depressed the number of

tilapias and *Synodontis*, a member of the catfish genus [39]. The author confirmed that thus, structural changes in the species composition of Lake Victoria's fish stocks may have been induced by the water hyacinth infestation of the lake

The Chinese grass carp is a fast growing fish which eats aquatic plants. It grows at a tremendous rate and reaches sizes of up to 32 kg [34]. It is an edible fish with a tasty white meat. It will eat submerged or floating plants and also bank grasses. The fish can be used for weed control and will eat up to 18 - 40% of its own body weight in a single day [30].

Other fish such as the tilapia, silver carp and the silver dollar fish are all aquatic and can be used to control aquatic weeds. The manatee or sea cow has also been suggested as another herbivore which could be used for aquatic weed control.

Water hyacinth has also been used indirectly to feed fish. Dehydrated water hyacinth has been added to the diet of channel catfish fingerlings to increase their growth [30]. It has also been noted that decay of water hyacinth after chemical control releases nutrients which promote the growth of phytoplankton with subsequent increases in fish yield [30].

Strategy for management of water Hyacinth

There are several popular control mechanisms for preventing the spread of, or eradication of, water hyacinth. Water hyacinth control methods fall into three main categories: physical, chemical and biological [39]. Its use as a livestock feed is considered as an effective physical control method. The best method of controlling water hyacinth is to prevent it from being introduced in to a marine and fresh water system. This can be done by educating the public about the problems that occur from disposal of unwanted water garden or aquarium plants in to marine and fresh water systems or by not properly cleaning boats, trailers, other water sports equipment, bait buckets, or fishing equipment to remove all plant material before moving the equipment to another fresh water system or within the lake boundaries itself.

Biological control is the most widely favored long term control method, being relatively easy to use, and arguably providing the only economic and sustainable control. Below we will briefly discuss each of these methods.

Mechanical control. Mechanical removal requires mechanical harvesters, self-tipping dump trucks, personnel (site supervisors, mechanics, and laborers), and disposal site for the removed weed biomass, push-boats, floating barriers, fuel and lubricants as the minimum items for implementation. Costs of removal per hectare are dependent on size of the infested area, distance to disposal site, density of the weed, accessibility of the areas infested and potential for re-infestation. It is not suitable for large infestations and is generally regarded as a short-term solution. For example Westerdahl and Getsinger [40] confirmed that mechanical controls such as harvesting have been used for nearly 100 years in Florida, but are ineffective for large scale control, very expensive, and cannot keep pace with the rapid plant growth in large water systems. The total cost per hectare is 681.3 USD [41] but costs of 250 USD per hectare have been reported. Manual/mechanical methods: For small ponds or lakes infested with water hyacinth, harvesting and removal of plant material from the water can be attempted. Care must be taken to remove all plant material, including small fragments. Removal of water hyacinth can be integrated with the preparation of organic fertilizer, so that it can add value for the community by preparing compost, silage for cattle feed and fish feed around each spots of removal sites. Effectiveness of manual control can be limited by wind movements, expansive mats covering large expanse area, location of infestation, water depth. In Lake Tana water hyacinth control efforts were implemented in a spirited way but there were challenges related namely; health threats of bilharzias, leech bites, malaria and cold water environment. Lack of appropriate equipment and protective wear and lack of support in initial stages were among the challenges [29].

Biological control. Biological control is the most widely favored long term control method, being relatively easy to use, and arguably providing the only economic and sustainable control. Three insects and a fungus have been extensively studied and subsequently released to control water hyacinth. According to Grodowitz, M.J. [42], these three insects have been released for the biological control of water hyacinth. These include two weevil species (*Neochetina* spp.) and a moth (*Sameodes alboguttalis*). Unfortunately large scale reductions in water hyacinth populations did not occur. Instead insect predation reduced plant height, decreased the number of seeds produced, and decreased the seasonal growth of the plants. This, in turn, allowed better boat access into plant mats, reduced use of herbicides, and resulted in less plant problems. In Louisiana, the seasonal growth of water hyacinth was reduced from a high of over 400,000 hectares per year to lows of only about 80,000 hectares. However this method

needs multidisciplinary study prior to introduce exotic species to water ecosystem that might cause for massive degradation of the resource. Grazing: most animals, except rabbits, do not readily eat the plant, possibly because its leaves are 95 percent water and have high tannin content. Rehabilitating some hotspot shores by wetland plants like papyrus is one of the remedy to overcome water hyacinth dominance. Papyrus serves as biological control. According to Anteneh [20] Survey report in Lake Tana the infestation water hyacinth was highest in the areas which indigenous Macrophytes such as papyrus devastated area. However area where papyrus is dominant, water hyacinth coverage becomes limited, because of Papyrus out competes water hyacinth.

Chemical control. Water hyacinth can be controlled using glyph sate as a foliar spray and copper complexes used only as a foliar spray. But herbicide use is more highly regulated in aquatic systems than in terrestrial systems. Chemical control, through the use of certain herbicides such as 2, 4-D or glyphosate, seems to be an economically feasible option in some countries, but not in others with less economic development. Westerdahl and Getsinger [40] report excellent control of water hyacinth by the use of the aquatic herbicides 2, 4-D or diquat. Chemical control is the least favored due the unknown long-term effects on the environment and the communities with which it comes into contact. In addition, in many countries public opinion is strongly against the use of chemicals in water, which is used for drinking purposes. So that cannot be recommended at this moment. Even though Manual removal requires a large labor force, and Governments of the developing world do not always have the means to pay for this operation, this would seems the best means of controlling the water hyacinth in Lake Tana.

CONCLUSION

Water hyacinth are negative environmental and economic impacts these include impairing and blocking of fishing grounds and fish breeding areas, reduced fish catches and fish quality, reduced tourism potential, reduced recession agricultural production and blockage of irrigation canals, loss of pasture lands and blocking of fishers movements could also be mentioned. Health impacts include increase in vector-borne diseases such as malaria, bilharzias and health related hazards from leeches and reptiles. Environmental impacts include deterioration in water quality, water loss through evapotranspiration and a decline of aquatic biodiversity. In conclusion, water hyacinth can be brought to make compost, mulching and to clean the sewage. It is a good way to change waste products into useful things. Apart from its ornamental value, the plant has been found useful as a source of animal feed as a source of fertilizers for use in agriculture, a source of bio- mass energy, a source of raw materials for building, handcraft making, paper and boards. In addition the plant has been found to be useful as a filter worth of solving man created problems of pollution in water bodies. More research is needed in order Water hyacinth has a multitude of negative consequences on economical, ecological and social. Current measurements were also made and they indicate that the wind is the main driving force of currents in the area. They also show that the circulation in the gulf is mainly horizontal rather than vertical. Water hyacinth could be an excellent source of proteins, vitamins, and minerals, and could be of particular value as a dietary supplement in countries where human diets are generally deficient in these nutrients. Water hyacinth control methods fall into three main categories: physical, chemical and biological. Its use as a livestock feed is considered as an effective physical control method.

Acknowledgments

The authors would like to thank the university of Gondar, College of Veterinary Medicine and Animal sciences for facilities and materials. I also pass my grateful for the other publishers who accessed the material to review this manuscript.

Competing interests

The authors declare that they have no conflict of interest with respect to the research, authorship or publications of this article.

REFERENCES

1. Cock M. 2001. Problems Caused by Water Hyacinth as an Invasive Alien Species. Case study from Wittenberg R, Cock MJW, eds, 2001. Invasive Alien Species: A Toolkit of Best Prevention and Management Practices. Wallingford, UK: CAB International, xvii-228.

2. Ogle BM, Dao HTA, Mulokozi G and Hambraeus L. 2001. Micronutrient composition and nutritional importance of gathered vegetables in Vietnam. *International Journal of Food Science and Nutrition*, 52(6): 485-499.
3. Wolverton B and McDonald RC. 1976. Don't waste waterweeds. *New Scientist*, 318-320.
4. Penfound WT, Earle TT. 1948. The Biology of the Water Hyacinth. *Ecological Monographs* 18: 447-472.
5. DeLoach CJ. 1976. Neochetina bruchi, a biological control agent of Water Hyacinth: host specificity in Argentina. *Ann. Entomol. Soc. Amer.* 69: 635-642.
6. Barrett SCH, Forno IW. 1982. Style morph distribution in New World populations of Eichhornia crassipes (Mart.) Solms-Laubach (water hyacinth). *Aquatic Botany*, 13: 299-306.
7. Bartodziej W, Weymouth G. 1995. Water bird abundance and activity on water hyacinth and Egeria in the St-Marks River, Florida. *Journal of Aquatic Plant Management*, 33: 19-22.
8. Brendonck L, Maes J, Rommens W, Dekeza N, Nihwatiwa T, Barson M, Callebaut V, Phiri C, Moreau K, Gratwicke B, Stevens M, Alyn N, Holsters E, Ollevier F, Marshall B. 2003. The impact of water hyacinth (Eichhornia crassipes) in a eutrophic subtropical impoundment (Lake Chivero, Zimbabwe). II. Species diversity. *Archiv Fur Hydrobiologie* 158: 389-405.
9. Lu JB, Wu JG, Fu ZH, Zhu L. 2007. Water hyacinth in China: A sustainability science based management framework. *Environmental Management* 40: 823-830.
10. Martinez Jimenez, M, Gomez Balandra MA. 2007. Integrated control of Eichhornia crassipes by using insects and plant pathogens in Mexico. *Crop Protection* 26: 1234-1238.
11. Twongo T, Bugenyi FWB, Wanda F. 1995. The potential for further proliferation of water hyacinth in lakes Victoria, Kyoga and Kwana and some urgent aspects for research. *Afr J Trop Hydrobiology Fish*, 6: 1-10.
12. Tackholm V, Drar M. 1950. Flora of Egypt II. Bull. Faculty Sci., Egypt Univ. Cairo 28: 441-448.
13. Mitchell, D.S., Pieterse, A.H., & Murphy, K.J. 1990. Aquatic weed problems and management in Africa. Pp.341-354 In A.H. Pieterse & K.J. Murphy, eds. *Aquatic Weed, the Ecology and Management of Nuisance Aquatic Vegetation*. Oxford, UK, Oxford University Press.
14. Ogutu-Ohwayo R, Hecky RE, Cohen AS, Kaufman L. 1997. Human impacts on the African Great Lakes. *Env Biol Fishs*, 50: 117-131.
15. Phiri G. 1997. An update on water hyacinth distribution and biological control. *Water Hyacinth Newsletter, CAB International* 6: 5.
16. Opande GO, Onyang JC, Wagai SO. 2004. Lake Victoria: The water hyacinth (Eichhornia crassipes (MART)), its socio-economic effects, control measures and resurgence in the Winam gulf. *Limnologica*, 34: 105-109
17. Kateregga E, Sterner T. 2007. Indicators for an invasive species: Water hyacinths Lake Victoria. *Ecological Indicators* 7; 362-370.
18. Food and Agricultural Organization, 1994. *Small water bodies and rivers in southern Africa*, edited by B. Marchal and M. Maes
19. Wondie A, Seid A, Molla E, Goshu G, G/kidan W/G, Dereje Tewabe A Sh, Genanew M. 2012. Preliminary Assessment of Water hyacinth (Eichornia crassipes) in Lake Tana. Proceedings of National Workshop (Biological Society of Ethiopia), Addis Ababa.
20. Anteneh W, 2015. Water Hyacinth Coverage Survey Report On Lake Tana Biosphere Reserve. Technical Survey Report Series 2.
21. Manson JG, Manson BE. 1958. Water hyacinth reproduces by seed in New Zealand. *New Zealand Jour. Agric.* 96: 191.
22. Matthews LJ, Manson BE, Coffey BT. 1977. Longevity of water hyacinth (Eichhornia crassipes (Mart.) Solms) seed in New Zealand. *Proc. 6th Asian-Pacific Weed Sci. Soc. Conf.*, Vol. 1: 263-267.
23. Schneider G. 1996. Rationale and Possible Strategies for Combating Aquatic weeds, UNEP Paper, Nairobi, Kenya.
24. Kateregga E, Sterner Th. 2008. Lake Victoria Fish Stocks and the Effects of Water Hyacinths on the Catch ability of Fish. Environment for Development discussion paper series.
25. Mailu AM. 2001. Preliminary Assessment of the Social, Economic, and Environmental Impacts of Water Hyacinth in the Lake Victoria Basin and the Status of Control. In *Biological and Integrated Control of Water Hyacinth, Eichhornia crassipes*, proceedings of the second meeting of the Global Working Group for the Biological and Integrated Control of Water Hyacinth, Beijing, China, 9–12 October 2000. Australian Centre for International Agricultural Research (ACIAR) Proceedings, no. 102: 130–139.
26. Henrylito D. Tacio, 2009. Water Hyacinth Ecological Value, Environmental Impacts. Report.
27. Makhanu KS. 1997. Impact of water hyacinth on Lake Victoria. *Water and Sanitation for All: Partnerships and Innovations. 23rd WEDC Conference.*
28. Saelthun, 1994. *Floating aquatic Macrophytes– Water hyacinths*
29. Edward R. 2013. *Lake Tana Water Hyacinth Management Strategy.*
30. Gopal B. 1987. *Water Hyacinth*. Elsevier, Amsterdam, Netherlands, p. 471.
31. Oyakawa N, Orlandi W, Valente EOL. 1970. The use of Eichhornia crassipes in the production of yeast feeds and forage. In *FAO Fisheries Technical Paper No. 187.*
32. Majid FZ. 1986. *Aquatic Weeds: Utility and Development*. 1st Ed India Agro Botanical Publishers.

33. Ogutu-Ohwayo R, Hecky RE, Cohen AS, Kaufman L. 1996. Human impacts on the African Great Lakes. *Environmental Biology of Fishes* 50: 117-131.
34. National Academy of Sciences. 1976. Making Aquatic Weeds Useful: Some Perspectives for Developing Countries.
35. Villadolid DV, Bunag DM. 1953. New uses for water hyacinths. *Philipp. Fish. Yearb.*, 80-1, 241-2
36. Sharma A. 1971. Eradication and Utilization of water hyacinth, A review. *Cur. Sci.* 40: 51-55.
37. Haider SZ. 1989. Recent Work in Bangladesh on the Utilization of Water Hyacinth, Intermediate Technology Development Group Commonwealth Science Council / Dhaka University.
38. Twongo T. 1998. Evolution of the Water hyacinth Problem in Uganda. Presidential Economic Council Report prepared for the Task Force on Water Hyacinth Control, May 1998.
39. Julien MH, Griffiths MW, Stanley JN. 2001. Biological control of water hyacinth 2. The moths *Niphograpta albiguttalis* and *Xubida infusellus*: biologies, host ranges, and rearing, releasing and monitoring techniques. Canberra, Australia. *ACIAR Monograph* No. 79.
40. Westerdahl, HE, Getsinger KD. 1988. Aquatic plant identification and herbicide use guide, volume II: Aquatic plants and susceptibility to herbicides. Technical report A-88-9. Department of the Army, Waterways Experiment Station, Corps of Engineers, Vicksburg, MS.
41. World Bank and Sida, 2012. Lake Victoria basin Water Hyacinth Surveillance, Monitoring and Control Strategy 2012 to 2030. East African Community, Lake Victoria Basin Commission Secretariat, P1 -145.
42. Grodowitz MJ 1998. An Active Approach to the Use of Insect Biological Control for the Management of Non-Naive Aquatic Plants. *Journal of Aquatic Plant Management.* 36:57-61.

Author Biography

The Authors have BSc degree in Animal Production and MSc in Fisheries and Wetlands Management. He was done his MSc Thesis title: "Diversity, Relative Abundance and Some aspects of the Biology of Fishes below Tisisat fall of the Blue Nile River. Ethiopia". Currently he had worked as lecturer at University of Gondar, College of Veterinary Medicine and Animal Sciences, research interests: Fisheries, Aquaculture, Aquatic Ecology, Climate change and Wetlands, Fish nutrition.