

Volume 8, No. 6, November 2018 * Bimonthly

ISSN: 2251-9939

Journal of Life Science and Biomedicine



Available online at www.jlsb.science-line.com

*Published by
Scienceline Publications*



Journal of Life Science and Biomedicine

(2251-9939)

J. Life Sci. Biomed. 8 (6): 90-100, November 25, 2018

Editorial Team

Editor-in-Chief: Parham Jabbarzadeh Kaboli

PhD of Molecular Biology and Cancer researcher; Faculty of Medicine and Health Sciences, University Putra, Malaysia ([Website](#); [Emails: researchgroups@drugdiscovery.ir](mailto:researchgroups@drugdiscovery.ir))

Managing Editor: Yusuf Kaya

PhD, Professor of Biology, Atatürk University, Erzurum, ([Website](#), [Email: ykaya@atauni.edu.tr](mailto:ykaya@atauni.edu.tr))

Executive Editor: Zohreh Yousefi

PhD candidate, Biosystematics, Atatürk University, Erzurum, Turkey ([Emails: zohreh.yousefi12@ogr.atauni.edu.tr](mailto:zohreh.yousefi12@ogr.atauni.edu.tr))

Language Editor: Samuel Stephen Oldershaw

Master of TESOL, The Humberston School & The Grimsby Institute, Nuns Corner, Grimsby, North East Lincolnshire, United Kingdom ([Email: s.s.oldershaw@hotmail.com](mailto:s.s.oldershaw@hotmail.com))

Associate Editors

Aleksandra K. Nowicka

PhD, Pediatrics and Cancer researcher; MD Anderson Cancer Center, Houston, Texas, USA ([Email: aknowicka@mdanderson.org](mailto:aknowicka@mdanderson.org))

Paola Roncada

PhD, Pharmacokinetics, Residues of mycotoxins in food and in foodproducing species, University of Bologna, Italy ([Email: paola.roncada@unibo.it](mailto:paola.roncada@unibo.it))

Tohid Vahdatpour

PhD, Assistant Prof., Physiology, Islamic Azad University, Iran ([Website](#); [Scopus](#); [Emails: vahdatpour@iaushab.ac.ir](mailto:vahdatpour@iaushab.ac.ir))

Veghar Hejazi

MD, Tabriz University of Medical Sciences, Tabriz, Iran ([Email: veggharhejazi@gmail.com](mailto:veggharhejazi@gmail.com))

Nefise Kandemir

MD, PhD, Department of Medical Genetics, Erciyes University, Kayseri, Turkey

Reviewers

Abolghasem Yousefi

PhD, Assistant Professor of Anesthesiology, Tehran University of Medical Sciences, Tehran, Iran ([Website](#); [Email: ayousefi@gmail.com](mailto:ayousefi@gmail.com))

Aleksandra K. Nowicka

PhD, Pediatrics and Cancer researcher; MD Anderson Cancer Center, Houston, Texas, USA ([Email: aknowicka@mdanderson.org](mailto:aknowicka@mdanderson.org))

Amany Abdin

PhD, Pharmacology; MSc, Medical Biochemistry; Tanta University, Egypt ([Emails: amanyabdin@med.tanta.edu.eg](mailto:amanyabdin@med.tanta.edu.eg), amanyahr@hotmail.com)

Babak Yousefi

Physician, General Surgery Resident at Hamedan University of Medical Science, Hamedan, Iran

Fazal Shirazi

PhD, Infectious Disease researcher at MD Anderson Cancer Center, Houston, Texas, USA

Fikret Çelebi

Professor of Veterinary Physiology; Atatürk University, Turkey ([Website](#); [Email: fcelebi@atauni.edu.tr](mailto:fcelebi@atauni.edu.tr))

Ghada Khalil Al Tajir

PhD, Pharmacology, Faculty of Medicine, UAE University, Al Ain, UAE

M.R. Ghavamnasiri

PhD, Professor of Oncology at Omid Cancer Hospital, MUMS; Cancer Research Center, Mashhad University of Medical Sciences, Iran

Kaviarasan Thanamegm

PhD of Marine Bioactive compounds, Department of Ecology and Environmental Sciences, Pondicherry University, India (Email: marinekavi@gmail.com)

Jahan Ara Khanam

PhD, Anti-cancer Drug Designer and Professor of UR; Department of Biochemistry and Molecular Biology, University of Rajshahi, Bangladesh

Mozafar Bagherzadeh Homaei

PhD, Plant Physiology, University of Isfahan, Isfahan, Iran

Osman Erganiş

Professor, PhD, Veterinary Microbiology, Selcuk University, Konya, Turkey

Paola Roncada

PhD, Pharmacokinetics, Residues of mycotoxins in food and in foodproducing species, University of Bologna, Italy (Email: paola.roncada@unibo.it)

Perumal Karthick

Professor, PhD, Marine Biology, Pondicherry University, Brookshabad Campus, Port Blair, Andamans. 744112, India (Email: karthickmicrobes@gmail.com)

Reza Khodarahmi

PhD, Biochemistry at KU; Pharmacy School, Kermanshah University, Kermanshah, Iran

Saeid Chekani Azar

PhD, Veterinary Physiology, Atatürk University, Erzurum, Turkey ([Google Scholar](#); [Emails: saeid.azar@atauni.edu.tr](mailto:saeid.azar@atauni.edu.tr); schekani@gmail.com)

Siamk Sandoughchian

PhD Student, Immunology, Juntendo University, Japan

Siva Sankar. R.

PhD, Marine Biology, Dept. of Ecology & Environmental Sciences, Pondicherry University, Puducherry - 605014, India (Email: sivauniverse@gmail.com)

Tohid Vahdatpour

PhD, Assistant Prof., Physiology, Islamic Azad University, Iran ([Website](#); [Scopus](#); [Google Scholar](#); [Emails: vahdatpour@iaushab.ac.ir](mailto:vahdatpour@iaushab.ac.ir))

Veghar Hejazi

MD, Tabriz University of Medical Sciences, Tabriz, Iran (Email: vegharhejazi@gmail.com)

Yusuf Kaya

PhD, Professor of Plant Biology, Atatürk University, Erzurum, Turkey (Email: ykaya@atauni.edu.tr)

Join JLSB Team

Journal of Life Sciences and Biomedicine (JLSB) as international journal is always striving to add diversity to our editorial board and operations staff. Applicants who have previous experience relevant to the position they are applying for may be considered for more senior positions (Section Editor) within JLSB. All other members must begin as Deputy Section Editors before progressing on to more senior roles. Editor and editorial board members do not receive any remuneration. These positions are voluntary.

If you are currently an undergraduate, M.Sc. or Ph.D. student at university and interested in working for JLSB, please fill out the application form below. Once your filled application form is submitted, the board will review your credentials and notify you within a week of an opportunity to membership in editorial board.

If you are PhD, assistant, associate editors, distinguished professor, scholars or publisher of a reputed university, please rank the mentioned positions in order of your preference. Please send us a copy of your resume (CV) or your [ORCID ID](#) or briefly discuss any leadership positions and other experiences you have had that are relevant to applied Medical and Pharmaceutical Researches or publications. This includes courses you have taken, editing, publishing, web design, layout design, and event planning. If you would like to represent the JLSB at your university, join our volunteer staff today! JLSB representatives assist students at their university to submit their work to the JLSB. You can also, registered as a member of JLSB for subsequent contacts by email and or invitation for a honorary reviewing articles.

Contact us at: editors@jlsb.science-line.com

Download [Application Form \(.doc\)](#)

Volume 8 (6); November 25, 2018**Research Paper****Acetylation Phenotype Impact on Early Postoperative Period in Viral Liver Cirrhosis.**

Ibadov RA, Omonov OA, and Ibragimov SKh.

J. Life Sci. Biomed., 8(6): 90-93, 2018;


pii:S225199391800014-8

Abstract

Objective. The aim of our study was to identify some pathogenetic mechanisms and unify prediction factors for the development of complications after portosystemic shunting. Material and Methods: The present research involved 45 patients with liver cirrhosis complicated by portal hypertension. Buccal swabs and spot urine samples were used to determine acetylation phenotypes. The genotype of each individual was determined by polymerase chain reaction. High-performance liquid chromatography was used to determine acetylation phenotypes. Results: Rapid acetylation was revealed in 7 patients (15.6%) and slow acetylation was found in 38 patients (84.4%). In slow acetylation phenotype, a considerable progression of liver cirrhosis was observed in comparison with rapid acetylators alanin aminotransferaz (ALT) on 74.4 % in slow acetylation phenotype (SACP) against 29.5% in rapid acetylation phenotype (RACP); total bilirubin on 111.8% in comparison with 42 %, respectively; the level of ammonia in blood was 247.8% compared to 62.5%). Recommendation: Taking into consideration the acetylation phenotype of liver cirrhosis patients can help in predicting possible side-effects and evaluate efficiency of drugs that are metabolized by N-acetylation.

Keywords: Acetylation Phenotype, Viral Liver Cirrhosis, Portal Hypertension, Central Portosystemic Shunting, Postoperative Period

[Full text-[PDF](#)] [[XML](#)]

**Research Paper****Water Hyacinth (*Eichhornia crassipes*) Biology and its Impacts on Ecosystem, Biodiversity, Economy and Human Well-being.**

Degaga AH.

J. Life Sci. Biomed., 8(6): 94-100, 2018;

pii:S225199391800015-8

Abstract

The aim of this review article was to show water hyacinth biology, chemical composition and its negative impacts on aquatic ecosystem, biodiversity, economy and human wellbeing. Water hyacinth is challenging the ecological stability of freshwater bodies. It is native to the Amazon Basin in Brazil and other nearby South American countries. In Africa, the first introduction of water hyacinth was in Egypt in 1880. In Ethiopia, water hyacinth was officially reported in 1956 in Koka Lake and the Awash River. Nutrients and temperature are considered the strongest determinants for water hyacinth growth and reproduction. Under favorable conditions, water hyacinths can double its mass every 5 days and it also grows from seed, which can remain viable for 20 years. Due to its extremely fast growth, the weed has become the major floating water weed of tropical and subtropical regions. In the absence of natural enemies, the weed quickly becomes invasive, colonizing slow moving waters resulting in thick and extensive mats which degrade aquatic ecosystems and limit their utilization. These mats affect fisheries and related commercial activities, functioning of irrigation canals, navigation, hydroelectric programmes and tourism. Its 95% mass weight is water from 5% dry matter 50% is silica and 30% is K, 15% N and 5% protein. The spread of this invasive plant is difficult to manage and not easy to reverse. Its impact is not only loss of biodiversity in aquatic ecosystems but also economic development and human wellbeing. It supports as breeding ground for vectors and pests. Hand removal is most effective for small infestations while mechanical harvesting can be an effective tool for removing larger infestations. The best method to control water hyacinth is to prevent it from entering a water body. This can be through education programs that have proved to be an effective tool in preventing further spread into catchments by people for ornamental purposes. So Ethiopian Government has to declare water hyacinth and other invasive species as a national pest and then put legislation in place to control them.

Keywords: Aquatic Ecosystem, Aquatic Weed, Invasive Plant Species, Fast Growth, Mat Formation

[Full text-[PDF](#)] [[XML](#)]



Journal of Life Science and Biomedicine



ISSN: 2251-9939

Frequency: Bimonthly

Current Issue: 2018, Vol: 8, Issue 6 (November)

Publisher: [SCIENCELINE](#)

The Journal of Life Science and Biomedicine is aimed to improve the quality and standard of life with emphasis on the related branches of science such as biology, physiology, biochemistry, zoology, anatomy, pathology and their applications and innovations in medicine and healthcare... [view full aims and scope](#)

<http://jlsb.science-line.com>

» JLSB indexed/covered by [NLM Catalog](#), [RiCeST \(ISC\)](#), [Ulrich's™](#), [SHERPA/RoMEO](#), [Genamics](#), [Google Scholar \(h-index= 10\)](#), [Index Copernicus](#), [ICV2015: 66.26...](#) [details](#)

» Open access full-text articles is available beginning with Volume 1, Issue 1.

» Full texts and XML articles are available in ISC-RiCeST.

» This journal is in compliance with [Budapest Open Access Initiative](#) and [International Committee of Medical Journal Editors' Recommendations](#).

ICMJE INTERNATIONAL COMMITTEE of
MEDICAL JOURNAL EDITORS

» High visibility of articles over the internet.

» Publisher Item Identifier [...details](#)

» This journal encourage the academic institutions in low-income countries to publish high quality scientific results, free of charges... [view Review/Decisions/Processing/Policy](#)



[ABOUT US](#)

| [CONTACT US](#)

| [PRIVACY POLICY](#)

Editorial Offices:

Atatürk University, Erzurum 25100, Turkey

University of Manitoba, Winnipeg, Manitoba R3T 2N2, Canada

University of Maragheh, East Azerbaijan, Maragheh 55136, Iran

Homepage: www.science-line.com

Phone: +98 914 420 7713 (Iran); +90 538 770 8824 (Turkey); +1 204 8982464 (Canada)

Emails:

administrator@science-line.com

saeid.azar@atauni.edu.tr

Acetylation Phenotype Impact on Early Postoperative Period in Viral Liver Cirrhosis

Ravshan Aliyevich IBADOV¹, Oybek Avazkhonovich OMONOV², and Sardor Khamdamovich IBRAGIMOV^{1*} 

¹Intensive Care Unit, Republican Specialized Scientific-Practical Medical Center of Surgery named after Academician V.Vakhidov, Tashkent, Uzbekistan

²Department of Portal Hypertension and Pancreatoduodenal Zone Surgery, Republican Specialized Scientific-Practical Medical Center of Surgery named after Academician V.Vakhidov, Tashkent, Uzbekistan

*Corresponding author's Email: dr.sardor.ibragimov@gmail.com

ABSTRACT

Objective. The aim of our study was to identify some pathogenetic mechanisms and unify prediction factors for the development of complications after portosystemic shunting. **Material and Methods:** The present research involved 45 patients with liver cirrhosis complicated by portal hypertension. Buccal swabs and spot urine samples were used to determine acetylation phenotypes. The genotype of each individual was determined by polymerase chain reaction. High-performance liquid chromatography was used to determine acetylation phenotypes. **Results:** Rapid acetylation was revealed in 7 patients (15.6%) and slow acetylation was found in 38 patients (84.4%). In slow acetylation phenotype, a considerable progression of liver cirrhosis was observed in comparison with rapid acetylators alanin aminotransferaz (ALT) on 74.4 % in slow acetylation phenotype (SAcP) against 29.5 % in rapid acetylation phenotype (RAcP); total bilirubin on 111.8 % in comparison with 42%, respectively; the level of ammonia in blood was 247.8% compared to 62.5%). **Recommendation:** Taking into consideration the acetylation phenotype of liver cirrhosis patients can help in predicting possible side-effects and evaluate efficiency of drugs that are metabolized by N-acetylation.

Original Article

PII: S225199391800014-8

Rec.	02 Nov	2018
Rev.	24 Nov	2018
Pub.	25 Nov	2018

Keywords

Acetylation Phenotype,
Viral Liver Cirrhosis,
Portal Hypertension,
Central Portosystemic
Shunting,
Postoperative Period

INTRODUCTION

Management of patients with liver cirrhosis complicated with portal hypertension after central portosystemic shunting. Hence, studying of acetylation polymorphism is currently relevant not only because many medical products are metabolized by acetylation reactions but also owing to better understanding the molecular basis of acetylation. These genetically caused metabolism variations of pharmaceuticals explain specific features of pharmacologic and therapeutic effect of drugs. Two genes found in humans are known to be responsible for activity of N-acetyl transferase. Recent research has shown that some alleles of these genes influence individual susceptibility to some diseases.

One of urgent problems in current pathologic physiology is studying the mechanisms of a disorder of detoxification functions of the liver in patients with various forms of liver pathology [1]. The hepatic endoplasmic network contains a family of isoenzymes of cytochrome P450 that is specific to various substrata. The processes of acetylation play an important part in interstitial metabolism. At present, acetylation phenotypes are considered to be a genetically determined ability of the body to metabolize compounds containing amino groups [2].

All pharmaceuticals pass the specific pharmacokinetic pathway by virtue of certain enzymes controlled genetically. Wide polymorphism in humans suggests that the fate of a pharmaceutical at any pharmacokinetic stage is associated with the polymorphic system of an enzyme or protein. It also causes diverse reactions of individuals to medicines [3].

To neutralize toxic products of metabolism or toxic substances in tissues some adaptable mechanisms, including those arranged in the toxigenic-kinetic, humoral, immunologic, and metabolic systems responsible for maintaining homeostasis in the body, have been developed in the course of evolution. Among them, the oxygen-dependent enzymes of the monooxygenase system play the important role [2]. Genetic differences in regulation, expression and activity of the genes, that code production of enzymes during the first and second

phases of xenobiotic biotransformation, can become a key factor of susceptibility to toxic effect of xenobiotics and development of a pathological process in the liver [4, 5].

Recently modern approaches of personalized medicine have been developed, e.g. assessment of the gene activity on the basis of studying the matrix RNA and drug metabolism [4]. Pharmacologic and kinetic research of pharmaceuticals is being conducted in many countries to evaluate the modes of drug dosing, considering individual variability of phenotypes of genetically determined biotransformation systems [5]. These studies will help not only to select the optimum doses of pharmaceuticals but to predict possible complications of the primary disease as well.

The aim of study was to identify some pathogenetic mechanisms and unify prediction factors for the development of complications after portosystemic shunting.

MATERIAL AND METHODS

Ethical approval

The review board and ethics committee of RSPMCS named after acad. V.Vakhidov approved the study protocol and informed consents were taken from all the participants.

The results of examination of 45 patients with viral liver cirrhosis complicated by portal hypertension (PH) have been analyzed. Morphological examination revealed large-nodule liver cirrhosis (LNLC) in over half of them (26 patients; 57.8%); 19 patients (42.2 %) had small-nodule liver cirrhosis (SNLC). In 39 patients, cirrhotic transformations of the liver were caused by viral hepatitis B, and in 6 patients it developed after viral hepatitis C. At the time of examination, antibodies to HCV were found in all 6 patients, and 39 patients had positive HBs-Ag. The patients were examined before and after central portosystemic shunting (PSS) with spleen preservation and after selective distal splenic-renal anastomosis (DSRA). The clinical course after the surgery was severe in 3 (6.7 %) patients, rather satisfactory in 5 (11.1 %) and uneventful in 37 (82.2 %) patients.

In addition to standard tests, the examination included evaluation of the level of reopirin metabolites, namely 4-amino-antipirin (4AAP) and N-acetyl-4-amino-antipirina (N-ac-4AAP) in urine. The latter method is specific because 4-AAP discharged with urine is a direct product of N- demethylation performed with microsomal monooxygenase system, while N-ac-4AAP is a product of further acetylation. The acetylating ability of the body was assessed by the method of Prebsting-Gavrilova modified by Anilova and Tolkachevsky. It was interpreted as slow if it did not reach 50%, and rapid when it made 50 % and more.

Before the surgery, a considerable decrease in excretion of reopirin metabolites was observed in all patients under study. For instance, in the SNLC patients, the level of 4 AAP in daily urine specimen was 3.6 times below the controls, and the level of the same metabolites in the LNLC patients was 7.36 times lower. The SNLC patients had 3-times lower N-ac-4AAP level, and that one in LNLC patients was 5.74 times lower. Rapid acetylation was revealed in 7 (15.6 %) patients, while slow acetylation was found in 38 patients (84.4 %).

RESULTS AND DISCUSSION

According to our findings, slow acetylation prevailed in patients with morphological variants of liver cirrhosis. For instance, the slow acetylation phenotype (SAcP) was found in 38 of 45 liver cirrhosis patients (84.4 %), while 7 (15.6 %) patients had the rapid acetylation phenotype (RAcP).

The comparative analysis of the basic blood biochemical parameters of patients with various types of acetylation made before and during the postoperative period has shown that an increase in the basic biochemical indicators of the liver did not depend on the type of acetylation. However, the values of these indicators were different in the compared groups. For instance, if a cytolytic component manifested itself as an increase in the levels of ALT and aspartate aminotransferase (AST) in blood of the patients before the surgery was almost identical in both groups, the postoperative indicator in the group of patients with RAcP was a little lower, than in the ones with SAcP.

The basic biochemical tests of blood before and after the postoperative period in patients with various morphological forms of cirrhosis demonstrated aggravation of these indicators depending on the liver cirrhosis form. As Figure 1 shows, a more favorable liver cirrhosis course in patients with rapid acetylation is obvious. For instance, if the ALT level increased from 212.7 ± 46.5 nmol/l to 383.4 ± 127.2 nmol/l in slow acetylation (i.e. a gain made 74.4 %), in the rapid type, the gain appeared to be considerably smaller: 29.5 % ($P < 0.05$). After surgery the total bilirubin level in the blood of patients with SAcP increased from 25.4 ± 6.7 to 53.8 ± 19.7 mcmmol/l that

made 111.8 %, while in the group with RAcP, hyperbilirubinemia was less expressed: before the surgery it was 23.1 ± 4.2 32.8 ± 8.1 $\mu\text{mol/l}$ or 42 % ($P < 0.01$) and after it.




Figure 1. Progression of the basic biochemical indicators of blood depending on the acetylation type in early postoperative period

When analyzing the total protein levels, it was revealed that in the patients with slow acetylation, the albumin fraction before the surgery had been 39.2 ± 2.9 g/L, while in the early postoperative period it had decreased to 32.8 ± 3.9 g/L. The ammonia level in the blood of patients with cirrhosis is rather demonstrative. The indicator before the surgery and in early postoperative periods again demonstrates the advantage of rapid acetylation. For instance, in the patients with slow acetylation, the ammonia level increased by 247.8 %, while in the patients with the rapid one, it increased by 62.5 % ($P < 0.01$).

The prothrombin time (PT) values before and after the surgery also differed, although to a lesser degree. In slow acetylation, PT decreased from 84.2 ± 6.8 to 75.4 ± 9.8 , (i.e. by 10.5 %), and in the rapid type, a decrease appeared to be considerably smaller: 7.9 % ($P < 0.5$).

Figure 2 presents the list and frequency of specific postoperative complications in the patients with different types of acetylation. The number of complications in patients with SAsP was observed to exceed the average incidence and specific complications developed more often than in rapid acetylators. Portosystemic encephalopathy was diagnosed in 6 patients and hepatic coma developed in 1 patient with SAcP, while in RAcP, only one patient had portosystemic encephalopathy of grades 1-2. Cholestasis was not observed in rapid acetylators, while in slow ones, it was observed in 2 cases. Edema and ascites developed in 7 patients with SAcP.

The correlation and comparative analysis demonstrated that parenchymatous-vascular decompensation in liver is characterized by: hepatic encephalopathy and mesenchymal and inflammatory response was observed in 36.8 % of patients with SAsP whereas in RAsP this complication developed only in one patient (14.3 %). No hemorrhage was observed in RAsP; in SAsP, it was found in 7.9 % of cases.




Figure 2. Progression of main biochemical indicators depending on the acetylation type

CONCLUSION

Slow acetylation phenotype mainly develops in liver cirrhosis patients (84.4 %), it being characterized by more often specific and nonspecific complications in the postoperative period irrespectively of the morphological form of cirrhosis. In slow acetylation, considerable liver cirrhotic progression in comparison with rapid acetylators was observed (ALT on 74.4 % in SAcP, against 29.5 % in RAcP, total bilirubin on 111.8 % compared to 42 %, the level of ammonia in blood was 247.8 % against 62.5 %, etc.).

Therefore, acetylation phenotypes of all patients with liver cirrhosis should be determined in the preoperative period since those ones with slow acetylation are at risk of possible specific and nonspecific complications in the postoperative period. Taking into consideration the acetylation phenotype of liver cirrhosis patients can help in predicting possible side-effects and evaluate efficiency of drugs that are metabolized by N-acetylation.

DECLARATIONS

Acknowledgements

This work was supported by Republican specialized scientific–practical medical center of surgery named after academician V.Vakhidov, Tashkent, Uzbekistan

Authors' Contributions

All authors contributed equally to this work.

Competing interests

The authors declare that they have no competing interests.

REFERENCES

1. Doll MA, Salazar-González RA, Bodduluri S, Hein DW. 2017. Arylamine N-acetyltransferase 2 genotype-dependent N-acetylation of isoniazid in cryopreserved human hepatocytes. *Acta Pharm Sin B*, 7(4):517-522.
2. Al-Ahmad MM, Amir N, Dhanasekaran S, John A, Abdulrazzaq YM, Ali BR, Bastaki S. 2017. Studies on N-Acetyltransferase (NAT2) Genotype Relationships in Emiratis: Confirmation of the Existence of Phenotype Variation among Slow Acetylators. *Ann Hum Genet*, 81(5):190-196.
3. Shin J, Kayser SR. Clinical pharmacy consultation for pharmacogenetic testing. 2009; 6(2):183-192.
4. Sychev DA, Ashraf GM, Svistunov AA, Maksimov ML, Tarasov VV, Chubarev VN, Otdelenov VA, Denisenko NP, Barreto GE, Aliev G. 2018. The cytochrome P450 isoenzyme and some new opportunities for the prediction of negative drug interaction in vivo. *Drug Des Devel Ther*, 12:1147-1156.
5. Verheijen RB. 2017. Clinical Pharmacokinetics and Pharmacodynamics of Pazopanib: Towards Optimized Dosing. *Clin Phrmacokinet*, 56(9): 987-997.

Water Hyacinth (*Eichhornia crassipes*) Biology and its Impacts on Ecosystem, Biodiversity, Economy and Human Well-being

Abera Hailu Degaga 

Department of Wildlife & Ecotourism Management, Wolkite University, P.O. Box 07, Wolkite, Ethiopia

Corresponding author's Email: aberabos@gmail.com; abera.hailu@wku.edu.et

ABSTRACT

The aim of this review article was to show water hyacinth biology, chemical composition and its negative impacts on aquatic ecosystem, biodiversity, economy and human wellbeing. Water hyacinth is challenging the ecological stability of freshwater bodies. It is native to the Amazon Basin in Brazil and other nearby South American countries. In Africa, the first introduction of water hyacinth was in Egypt in 1880. In Ethiopia, water hyacinth was officially reported in 1956 in Koka Lake and the Awash River. Nutrients and temperature are considered the strongest determinants for water hyacinth growth and reproduction. Under favorable conditions, water hyacinths can double its mass every 5 days and it also grows from seed, which can remain viable for 20 years. Due to its extremely fast growth, the weed has become the major floating water weed of tropical and subtropical regions. In the absence of natural enemies, the weed quickly becomes invasive, colonizing slow moving waters resulting in thick and extensive mats which degrade aquatic ecosystems and limit their utilization. These mats affect fisheries and related commercial activities, functioning of irrigation canals, navigation, hydroelectric programmes and tourism. Its 95% mass weight is water from 5% dry matter 50% is silica and 30% is K, 15% N and 5% protein. The spread of this invasive plant is difficult to manage and not easy to reverse. Its impact is not only loss of biodiversity in aquatic ecosystems but also economic development and human wellbeing. It supports as breeding ground for vectors and pests. Hand removal is most effective for small infestations while mechanical harvesting can be an effective tool for removing larger infestations. The best method to control water hyacinth is to prevent it from entering a water body. This can be through education programs that have proved to be an effective tool in preventing further spread into catchments by people for ornamental purposes. So Ethiopian Government has to declare water hyacinth and other invasive species as a national pest and then put legislation in place to control them.

Original Article

PII: S225199391800015-8

Rec.	28 Sep	2018
Acc.	19 Oct	2018
Rev.	15 Nov	2018
Pub.	25 Nov	2018

Keywords

Aquatic Ecosystem,
Aquatic Weed,
Invasive Plant Species,
Fast Growth,
Mat Formation

INTRODUCTION

The spread of invasive species is difficult to manage and not easy to reverse, this threatens not only biodiversity of aquatic ecosystems but also economic development and human wellbeing [1]. Water hyacinth (*Eichhornia crassipes*) is an invasive aquatic plant associated with a variety of ecological and economic effects on freshwater ecosystems [2]. It is a free-floating aquatic plant that grows in ponds or slow moving waterways. It is a perennial monocotyledonous crop that belongs to the Pontederiaceae family. It is native to the Amazon Basin in Brazil and other nearby South American countries [3]. And Holm, *et al.*, [4] reported that, *E. crassipes*, a native of South America, is a major freshwater weed in most of the frost-free regions of the world and is generally regarded as the most troublesome aquatic plant. It is considered the worst aquatic weed in the world [5]. In Africa, the first introduction of water hyacinth was in Egypt in 1880 [6]; the main aquatic weed in East Africa is Water hyacinth [5]. In Ethiopia, water hyacinth was officially reported in 1965 in Koka Lake and the Awash River [7, 8] and infestation of Lake Tana was officially recognized in 2011 [9]. It has been recognized as the most damaging aquatic weed in Ethiopia since 1965 [8]. In Lake Tana in addition to water hyacinth, other two floating invasive weeds: Azolla and Water Lettuce, were reported [10].

Water hyacinth reproduces both sexually and asexually. The rapid increase and spread of the plant into new areas is due particularly to its vegetative reproduction, a single plant being able to develop very rapidly a significant infestation [11]. Water hyacinth has a rapid propagation and morphological characteristics that makes the weed well adapted to rapid distance dispersal and successful colonization of varying habitats in a

short time [12]. Moving easily with water currents, winds or other accidental means, such as fishing nets and boats, the plant invaded rivers, canals, ponds, lakes, dams and other freshwater bodies. In the absence of natural enemies, the weed quickly becomes invasive, colonizing slow moving waters resulting in thick and extensive mats [13] which degrade aquatic ecosystems and limit their utilization [14]. The negative impacts of water hyacinth are due to its dense, impenetrable mats which restrict access to water. These mats affect fisheries and related commercial activities, functioning of irrigation canals, navigation/transport, hydroelectric programmes and tourism [15].

Nutrients and temperature are considered the strongest determinants for water hyacinth growth and reproduction [17]. Salinity constraints generally limit water hyacinth establishment in coastal areas and within estuaries [17]. Due to its extremely fast growth, the weed has become the major floating water weed of tropical and subtropical regions. There for the aim of this review was to show water hyacinth biology, chemical composition and its negative impacts on aquatic ecosystem biodiversity, economy and human wellbeing. And to show water hyacinth is challenging the ecological stability of freshwater ecosystems.

Biology, chemical composition and ecology of water hyacinth

The *E. crassipes* growth is extremely rapid and forms large populations of inter-connected shoots which is impenetrable mat. It forms dense, interlocking mats due to its rapid reproductive rate and complex root structure [18]. The flowers are bluish purple, large and self-fertile. The seeds are produced in large numbers and are contained in capsules, each capsule containing up to 300 seeds [19]. The seeds can remain viable for 5-20 years [20]. The plant can also reproduce vegetative through the production of horizontal stolons. Rakotoarisoa, *et al.*, [11] described that due to its high reproduction rate, the complex root structure and the formation of dense mats with up to two million plants per hectare can be found. Under favorable conditions, water hyacinth can double its mass every 5 days and it also grows from seed, which can remain viable for 20 years or longer [21, 22]. The biotic seeds dispersals are birds thought to be transported over long distances (e.g. waterfowl and shore birds) and if coated in mud they may cling to both mammals and birds [23, 24]. While, wind is the abiotic dispersal, it will readily move the plant and the upright leaves in lakes and canals. Along rivers, water flow is the prime mover of vegetative material but strong winds may sometimes blow the plant upstream.

Water hyacinth draws all its nutrients directly from water. It absorbs heavy metals [25], organic contaminants [26], and nutrients from the water column [27]. It comprises 95% water and 5% dry matter of which 50% is silica, 30% Potassium, 15% Nitrogen and 5% protein [28]. While Roger and Davis [29] reported that the uptake of nitrogen by water hyacinth is 5 to 10 times as rapidly as phosphorous. It has been known to thrive well in nutrient-enriched fresh waters in tropical climatic zones. For this purpose it has been used in wastewater treatment facilities [30].

The structure of a macrophyte assemblage plays a large role in determining composition of phytoplankton, zooplankton, fish, and birds in freshwater ecosystems [31]. A shift in the primary-production base of a lake can resonate throughout the ecosystem, affecting multiple trophic levels both directly through changes in habitat availability and indirectly through shifts in energy pathways. Free floating plants are able to monopolize light and absorb nutrients from the water column, preventing phytoplankton and submersed vegetation from obtaining sufficient resources for photosynthesis [32]. Altering ecosystem services and processes, reducing native species abundance and richness, and decreasing genetic diversity of ecosystems [33] and also water hyacinth affects diversity, distribution and abundance of life in aquatic environments [34].

Water hyacinth impacts on biodiversity loss

Water hyacinth is challenging the ecological stability of freshwater bodies [35], out-competing all other species growing in the vicinity, posing a threat to aquatic biodiversity [22]. Besides suppressing the growth of native plants and negatively affecting microbes, water hyacinth prevents the growth and abundance of phytoplankton under large mats, ultimately affecting fisheries [36]. This is because fish feed on phytoplankton. According to the Millennium Ecosystem Assessment [37], freshwater ecosystems are among the most significantly human-altered systems in the world. While invasive species are considered the leading threat to global aquatic biodiversity [38].

Most water hyacinth effects are lower phytoplankton productivity and dissolved oxygen concentrations beneath mats [39, 40]. Reduced phytoplankton productivity can decrease zooplankton abundance by decreasing food availability [19, 41]. It also affects diversity, distribution and abundance of life in aquatic environments and enhances evapo-transpiration, thus affecting all aquatic organisms. The death and decay of water hyacinth

vegetation in large masses create anaerobic conditions and production of lethal gases [34]. Coverage of water hyacinth causes de-oxygenation of water, and at times anoxia below the dense mats [42]. Water quality effects include higher sedimentation rates within the plant's complex root structure and higher evapo-transpiration rates from water hyacinth leaves when compared to evaporation rates from open water [43]. A shift in the primary-production base of a lake can resonate throughout the ecosystem, affecting multiple trophic levels both directly through changes in habitat availability and indirectly through shifts in energy pathways [44]. Dissolved oxygen less than 5 mg per liter are known to adversely affect function and survival of most fishes and less than 2 mg per liter can lead to fish kills [45]. This low level of dissolved oxygen along the water column will happen when large water hyacinth mats prevent light infiltration or when a relatively large area of plants decompose at the same time.

In the 1950s, within three years of its first sighting, it had spread 1600 km along the Congo River [24]; in Lake Tana in 2012 after a year of infestation officially recognized its estimation coverage was 20,000 hectares [46]. This shows that if once introduced to favorable habitats, especially open waters, *E. crassipes* may spread very rapidly and can form dense monotypic mats. Which is impenetrable and blocking access both for transportation and aquatic living organisms in addition to adding to much organic matter to water bodies leading anoxia condition. The introduction and spread of non-native species contribute to the loss of aquatic species [47].

Economic impacts of water hyacinth

Because of its rapid growth and mat formation, *E. crassipes* has a range of detrimental effects on the economy sector. The dense mats disrupt socioeconomic and subsistence activities for example ship and boat navigation, restricted access to water for recreation, fisheries, and tourism [22, 48]. It physically interferes with water transport, communication and access. Infestations are increasing in Ethiopia, creating a range of problems including restricted access [49]. In Nigeria, Alimi and Akinyemiju [50] showed that costs of fuel and repairs to boats on infested waterways was approximately three times that on un-infested waterways. Economic losses also result from interference with recreational uses of water bodies [43, 49]. Heavy infestations by Water Hyacinth make fishing very difficult, or impossible [5]. Fishermen are being troubled by a reduced range of fish species, loss of nets and impeded access [51]. Water hyacinth was perceived to affect fisheries through reduced levels of production, a reduction in species diversity, poor quality fish, rising cost of operation resulting in lower income to fishers and higher prices to consumers [52].

In Lake Victoria mats blocked breeding, nursery, and feeding grounds for economically important fish species, such as tilapia and Nile perch. Because water hyacinth mats can reduce natural predation and fisheries catchability, leading to increased abundance of certain species [53]; but mats can also exclude certain species from important breeding, nursery, and feeding grounds [54]. Expensive barriers or mechanical damage to hydro-electric installations and other structures such as bridges; for example, to the Owen Falls Dam on Lake Victoria [55], there are also similar concerns in South Africa [56], and Ethiopia [27].



Figure 1. Shore of Lake Tana (a) Cattle grazing and water hyacinth (b) Hippopotamus select grass but do not graze water hyacinth [10]

Water hyacinth has limited beneficial uses. Local communities around Lake Tana, they are worried about the invasion of their shore farm and grazing lands and cattle grazing the water hyacinth when there is no grass [10]. It cannot be used as a livestock feed because it contains too much silica, calcium oxalate, potassium and too little protein [30]. Therefore, this leads reducing grazing potential, have a negative impacts on animal health, milk and meat quality (Figure 1) and economical reduction on the livestock sector of the country. Dereje, *et al.*, [34] mentioned that expansion of water hyacinth around Lake Tana and its competition with the native species

the submerging grasses and other native species becomes devastated. These affect a lot of cattle which are directly and indirectly dependent on the grass. And also shore area floras which would be important for fish breeding grounds and livestock forage source in the vicinity become damaged.

Water hyacinth impacts on human wellbeing

E. crassipes may reduce water quality in various ways and encourage mosquitoes, snails and other organisms associated with human illnesses, including malaria, schistosomiasis, encephalitis, filariasis and cholera [43, 56] It also increased health hazards i.e. incidence of malaria and schistosomiasis [15]. FAO [5] also described that, this weed represents an environmental problem as well and indirectly a public health problem, since it may create a microhabitat suitable for the breeding of many vectors of human diseases and for hosting poisonous snakes. The infestation of water hyacinths (*Eichhornia crassipes*) in African lakes has increased breeding site availability for malaria vector species, *An. funestus* complex were reported from a water hyacinth mat [57]. Malaria vectors are able to breed amongst water hyacinth mats in Lake Victoria [58].

Water hyacinth control methods

Water hyacinth is extremely difficult to eradicate once established, the goal of most management efforts is to minimize economic costs and ecological change [44] The optimum control method depends on the specific conditions of each affected location such as the extent of water hyacinth infestation, regional climate, and proximity to human and wildlife [36]. Hand removal is most effective for small infestations such as small dams and drains. It is highly laborious and should only be used where the rate of removal can exceed the rate of re-growth. It should be done before flowering and seed set in spring [1]. Mechanical shredding of water hyacinth is cheaper than harvesting [59], but there are significant consequences of allowing the plant to die and decompose within the system. Understanding the density threshold at which water hyacinth starts to impact ecosystems, society, and local economies will help management establishment goals for population control that can maximize the social benefits while minimizing the costs of the invasive species [44]. Therefore, the best method to control water hyacinth is to prevent it from entering a water body.

Biological control is most effective on larger infestations but it can take several years for it to provide successful control. It involves the use of natural enemies including plant pathogens [36, 60]. The aim of any biological control is not to eradicate the weed, but to reduce its abundance to a level where it is no longer problematic. While there exists several native enemies of water hyacinth, two South American weevil beetles (*Neochetina eichhorniae* and *Neochetina bruchi*) and two water hyacinth moth species (*Niphograpta albiguttalis* and *Xubida infusella*) have had effective long-term control of water hyacinth in many countries, notably at Lake Chivero [28], Lake Victoria (Kenya), Louisiana (USA), Mexico, Papua New Guinea and Benin [35, 60, 61, 62]. Researchers have identified another tiny insect, *Megamelus scutellaris*, from South America which is highly host-specific to water hyacinth and does not pose a threat to native or economically important species [63].

CONCLUSION AND RECCOMENDATION

Biological alien invasions are a major driver of biodiversity loss worldwide. Water hyacinth (*Eichhornia crassipes*) is common and widely distributed all over the world, is challenging the ecological stability of freshwater ecosystems. The spread of invasive alien species is neither easy to manage nor easy to reverse. They are threatening not only biodiversity but also economic development and human wellbeing. Threats are destruction of biodiversity; oxygen depletion and reduced water quality; breeding ground for pests and vectors; blockage of waterways hampering agriculture, fisheries, recreation and hydropower; fishing, grazing and other agricultural activities by forming impenetrable thickets and hindering movements of humans and animals, and destroying and replacing natural biodiversity. Proliferation of water hyacinth is a symptom of broader watershed management and pollution problems.

The best method to control water hyacinth is to prevent it from entering a water body. Development of national and local policies for the detection, control and eradication of invasive species within and around aquatic ecosystems, farm lands, communal lands and in all ecosystems is required to prevent impacts of invasive species ahead not only on biodiversity loss but also, ecosystem and economy of a country. Therefore, the recommendation based on this review is that Ethiopian Government has to declare water hyacinth and other invasive species as a national pest and then put legislation in place to control them. Since Ethiopia being a member of Convention on Biological Diversity (CBD) which urges the parties to "prevent the introduction of,

control, or eradicate those alien species which threaten ecosystem, habitat or species"; the impact of invasive weeds on environment, article 8(h) of the CBD signed by 161 countries at the Earth Summit [64] .

DECLARATIONS

Acknowledgements

This review paper was presented in Ethiopian Fishery and aquatics science association during this time I took comments and suggestions. So the author will acknowledge the association members and scholars for their valuable comments.

Competing interests

The author declare that there is no any competing interests.

REFERENCES

1. UNEP. 2012. Fifth Global Environment Outlook (GEO5): Environment for the future we want. United Nations Environment Programme, Nairobi.
2. Center TD, (ed). 1994. Biological Control of weeds: water hyacinth and water lettuce. Intercept, Andover.
3. Chunkao K, Nimpee C and Duangmal 2012. The King's initiatives using water hyacinth to remove heavy metals and plant nutrients from wastewater through Bueng Makkasan in Bangkok, Thailand. *Ecological Engineering*; 39: 40–52.
4. Holm L, Doll J, Holm E, Pancho J and Herberger J. 1997. World Weeds. Natural Histories and Distribution. New York, USA: John Wiley and Sons, Inc
5. FAO. 2002. Management of problematic aquatic weeds in Africa FAO efforts and achievements during the period 1991-2001, Rome, Italy
6. Warnimont FJ. 1965. Problem in the Congo. The water hyacinth problem in the Congo catchment area. Democratic Republic of Congo, Leopoldville; 23.
7. Firehun Y, Struik PC, Lantinga EA and Taye T. 2014. Water hyacinth in the rift valley water bodies of Ethiopia its distribution socioeconomic importance and management. *IJCAR*; 3: 67-75.
8. Wondie Z. 2013. Assessment of water Hyacinth (*Eichhornia crassipes* (Mart) Solms) in relation to water quality, composition and abundance of plankton and macro-invertebrates in the north-eastern part of Lake Tana, Ethiopia.
9. Ayalew W, Ali S, Eyayu M, Goraw G, W/Gebriel G/K, Agegnehu Sh, Dereje T and Muluneh G. 2012. Preliminary Assessment of Water hyacinth (*Eichhornia crassipes*) in Lake Tana. Proceedings of National Workshop (Biological Society of Ethiopia), Addis Ababa
10. Wassie A, Minwuyelet M, Ayalew W, Dereje T, Woldegebrael W/K, Addisalem A and Wondie E. 2014. Water hyacinth coverage survey report on Lake Tana, Technical Report Series 1
11. Rakotoarisoa TF, Waeber PO, Richter T and Mantilla Contreras IJ. 2015. Water hyacinth (*Eichhornia crassipes*) any opportunities for the Alaotra wetlands and livelihoods. *MCD*; 10: 128-136.
12. Obeid M. 1984. Water hyacinth, (*Eichhornia crassipes* Mart.) solms. In Sudan. Proceedings of the international conference on water hyacinth. Hyderabad, India, pp. 145-148.
13. Edwards D and Musil CJ. 1975. *Eichhornia crassipes* in South Africa – a general review. *J. Limnological Soc. Southern Afr.*; 1: 23-27.
14. Hill MP and Coetzee JA. 2008. Integrated control of water hyacinth in Africa. *EPPO Bull.* 38: 452-457
15. Navarro L and Phiri G. 2000. Water hyacinth in Africa and the Middle East. A survey of problems and solutions. International Development Research Centre, Ottawa (CA).
16. Wilson JRU, Ajuonu O, Center TD, Hill MP, Julien MH, Katagira FF, Neuenschwander P, Njoka SW, Ogwang J, Reeder RH and Van T. 2007. The decline of water hyacinth on Lake Victoria was due to biological control by *Neochetina* spp. *Aquatic Botany*; 87: 90-93.
17. Mangas-Ramirez E and Elias-Gutierrez M. 2004. Effect of mechanical removal of water hyacinth (*Eichhornia crassipes*) on the water quality and biological communities in a Mexican reservoir. *Journal of Aquatic Health and Management*; 7: 161-168.
18. Mitchell DS. 1985. Surface-floating aquatic macrophytes. pp. 109-124 In: The Ecology and Management of African Wetland Vegetation P. Denny, editor. W. Junk Publishers, Dordrecht.
19. Maceina MJ, Cichra M, Betsill R and Bettoli P. 1992. Limnological changes in a large reservoir following vegetation removal by grass carp. *Journal of Freshwater Ecology*; 7:81-93.
20. Manson JG and Manson BE. 1958. Water hyacinth reproduces by seed in New Zealand. *New Zealand Jour. Agric.*; 96: 191
21. Frezina NCA. 2013. Assessment and utilization of water hyacinth in the water bodies of Tamil Nadu. *IJSRP*; 2: 58-77.
22. Patel S. 2012. Threats, management and envisaged utilizations of aquatic weed *Eichhornia crassipes*: an overview. *Rev Environ Sci Biotechnol*; 11:249–259.
23. Batcher MS. 2000. *Eichhornia crassipes* (Martius) Solms. Element Stewardship Abstract. Arlington, USA: The Nature Conservancy.

24. Holm LG, Weldon LG and Blackburn RD. 1969. Aquatic weeds. *Science*; 166:699-709.
25. Tiwari S, Dixit S and Verma N. 2007. An effective means of biofiltration of heavy metal contaminated water bodies using aquatic weed *Eichhornia crassipes*. *Environmental Monitoring and Assessment*; 129: 253-256.
26. Zimmels Y, Kirzhner F and Malkovskaja A. 2007. Advanced extraction and lower bounds for removal of pollutants from wastewater by water plants. *Water Environment Research*; 79: 287-296.
27. Aoi T and Hayashi T. 1996. Nutrient removal by water lettuce (*Pistia stratiotes*). *Water Science and Technology*; 34: 407-412.
28. Makhanu KS. 1997. Impact of Water hyacinth in Lake Victoria. In: Water and Sanitation for all: Partnerships and Innovations. 23rd Water Engineering and Development Centre Conference Durban, South Africa.
29. Roger HH and Davis DE. 1972. Nutrient removal by water hyacinth. *Weed Sci.*; 20: 423-427; In: Dereje T, Erkie A, Wondie Z and Brehan M. 2017. Identification of impacts, some biology of water hyacinth (*Eichhornia crassipes*) and its management options in Lake Tana, Ethiopia. *Net Journal of Agricultural Science*; 5(1): 8-15.
30. Osumo MW. 2001. Effects of water hyacinth on water quality of winam gulf, Lake Victoria. UNU-Fisheries Training Programme, Skulagata 4 120 Reykjavik, Iceland
31. Meerhoff M, Fosalba C, Bruzzone C, Mazzeo N, Noordoven W and Jeppesen E. 2006. An experimental study of habitat choice by *Daphnia*: plants signal danger more than refuge in subtropical lakes. *Freshwater Biology*; 51: 1320-1330.
32. McVea C and Boyd CE. 1975. Effects of water-hyacinth cover on water chemistry, phytoplankton, and fish in Ponds. *Journal of Environmental Quality*; 4: 375-378.
33. Rands M, Adams W, Bennun L, Butchart S, Clements A, Coomes D, Entwistle A, Hodge I, Kapos V, Scharlemann J, Sutherland W and Vira B. 2010. Biodiversity conservation: Challenges beyond 2010. *Science*; 329: 1298-1303.
34. Dereje T, Erkie A, Wondie Z and Brehan M, 2017. Identification of impacts, some biology of water hyacinth (*Eichhornia crassipes*) and its management options in Lake Tana, Ethiopia. *Net Journal of Agricultural Science*; 5(1): 8-15.
35. Gichuki J, Omondi R, Boera P, Tom Okorut T, SaidMatano A, Jembe T and Ofulla A, 2012. Water Hyacinth *Eichhornia crassipes* (Mart.) Solms-Laubach Dynamics and Succession in the Nyanza Gulf of Lake Victoria (East Africa): Implications for Water Quality and Biodiversity Conservation. *The Scientific World Journal*,
36. Villamagna A and Murphy B. 2010. Ecological and socio-economic impacts of invasive water hyacinth (*Eichhornia crassipes*): a review. *Freshwater Biology*; 55: 282- 298
37. Millennium Ecosystem Assessment (2005). *Ecosystems and Human Well-being: Biodiversity Synthesis*. World Resources Institute, Washington, DC
38. Mack RN, Simberloff D, Lonsdale WM, Evans H, Clout M and Bazzaz FA. 2000. Biotic invasions: Causes, epidemiology, global consequences, and control. *Ecological Applications* 10: 689-710.
39. Mironga JM, Mathooko JM and Onywere SM. 2011. The Effect of Water Hyacinth (*Eichhornia Crassipes*) Infestation on Phytoplankton Productivity in Lake Naivasha and the Status of Control. *Journal of Environmental Science and Engineering*; 5(10): 1252-1261
40. Rommens W, Maes J, Dekeza N, Inghelbrecht P, Nhiwatiwa T, Holsters E, Ollevier F, Marshall B and Brendonck L. 2003. The impact of water hyacinth (*E. crassipes*) in a eutrophic subtropical impoundment (Lake Chivero, Zimbabwe). I. Water quality. *Archiv Fur Hydrobiologie*; 158: 373-388
41. Richards DI, Small J and Osborne J. 1985. Response of zooplankton to the reduction and elimination of submerged vegetation by grass carp and herbicides in four Florida lakes. *Hydrobiologia*; 123: 97-108.
42. Gerry H, Waage J and Phiri G. 1997. The problem in tropical Africa. Report prepared for the first meeting on and international water hyacinth consortium held at the World Bank, Washington 18-19 March, 1997.
43. Gopal P, 1987. *Aquatic Plant studies 1: Water hyacinth*. Netherlands Elsevier Science Publishers B. V
44. Amy MV. 2009. Ecological effects of water hyacinth (*Eichhornia crassipes*) on Lake Chapala, Mexico. Dissertation submitted to the Faculty of the Virginia Polytechnic Institute and State University, Blacksburg, Virginia.
45. Chapman D, (ed). 1996. *Water quality assessments: A guide to the use of biota, sediments and water in environmental modeling*. Chapman & Hall, London.
46. Bureau of Environmental Protection, Administration and Use (BoEPLAU). 2012. In: Wassie A, Minwuyelet M, Ayalew W, Dereje T, Woldegebrael W/K, Addisalem A and Wondie E. 2014. Water hyacinth coverage survey report on ake Tana, Technical Report Series 1
47. Suski CD and Cooke SJ, 2007. Conservation of aquatic resources through the use of freshwater protected areas: Opportunities and challenges. *Biodiversity and Conservation*; 16: 2015-2029.
48. Ndimele P, Kumolu-Johnson C and Anetekhai M. 2011. The invasive aquatic macrophyte, water hyacinth {*Eichhornia crassipes* (Mart.) Solm-Laubach: Pontedericeae}: problems and prospects. *Res J Environ Sci* 5:509-520.
49. Aweke G. 1994. The water hyacinth (*Eichhornia crassipes*) in Ethiopia. *Bulletin des Séances, Académie Royale des Sciences d'Outre-Mer*; 39(3):399-404.
50. Alimi T and Akinyemiju OA. 1991. Effects of water hyacinth on water transportation in Nigeria. *Journal of Aquatic Plant Management*; 29:109-112.
51. Terry PJ, 1996. The water hyacinth problem in Malawi and foreseen methods of control. *Strategies for Water Hyacinth Control*. Report of a panel of experts meeting, 1995, Fort Lauderdale, USA. Rome, Italy: FAO, 59-81.
52. LVEMP. 1995. Lake Victoria Environmental Management programme. Report submitted by Kenya, Uganda and Tanzania to World Bank.

53. Kateregga E and Sterner T. 2009. Lake Victoria fish stocks and the effects of water hyacinth. *The Journal of Environment & Development*; 18: 62-78.
54. Twongo T and Howard G. 1998. Ways with weeds. *New Scientist*; 159: 57-57
55. Hill G, Cock M and Howard G. 1999. A global review of water hyacinth - its control and utilization. CABI Bioscience and IUCN
56. Harley KLS, Julien MH and Wright AD. 1996. Water hyacinth: a tropical worldwide problem and methods for its control. In: Julissa RS. & Pedro A-R., 2013. *Eichhornia crassipes* (water hyacinth). Department of Botany-Smithsonian NMNH, Washington DC, USA, /accessed on Dec. 25, 2017/.
57. Minakawa N, Seda P and Yan G. 2002. Influence of host and larval habitat distribution on the abundance of African malaria vectors in western Kenya. *Am J Trop Med Hyg*; 67:32–38.
58. Noboru M, Gabriel OD, George OS, Kyoko F and Sammy MN. 2012. Malaria Vectors in Lake Victoria and Adjacent Habitats in Western Kenya. *PLoS One*; 7(3): e32725
59. Greenfield BK, Blankinship M and McNabb TJ. 2006. Control costs, operation, and permitting issues for non-chemical plant control: Case studies in the San Francisco Bay-Delta Region, California. *Journal of Aquatic Plant Management*; 44: 40-49.
60. Dagno K, Lahlali R, Diourte M and Haissam J, 2012. Fungi occurring on water hyacinth (*Eichhornia crassipes* [Martius] Solms-Laubach) in Niger River in Mali and their evaluation as Mycoherbicides. *J. Aquat. Plant Manage*; 50: 25-32.
61. Venter N, Hill M, Hutchinson S and Ripley B. 2012. Weevil borne microbes contribute as much to the reduction of photosynthesis in water hyacinth as does herbivory. *Biological Control*; 64: 138–142.
62. Williams A, Hecky R and Duthie H. 2007. Water hyacinth decline across Lake Victoria-Was it caused by climatic perturbation or biological control? A reply. *Aquatic Bot*; 87:94–96.
63. Coetzee J, Hill M, Julien M, Center T and Cordo H. 2009. *Eichhornia crassipes* (Mart.) Solms–Laub. (Pontederiaceae).
64. Convention on Biological Diversity (CBD). 1992. United Nation (UN).

Instructions for Authors

Manuscript as Original Research Paper, Review and Case Reports are invited for rapid peer-review publishing in the *Journal of Life Science and Biomedicine*. Considered subject areas include: Biocontrol, Biochemistry, Biotechnology, Bioengineering, Neurobiology... [view full aims and scope](#)

[JLSB EndNote Style](#)

[Manuscript Template \(.doc\)](#)

[Sample Articles](#)

[Declaration form](#)

[Policies and Publication Ethics](#)

Submission

The manuscript and other correspondence should preferentially be [submit online](#). Please embed all figures and tables in the manuscript to become one single file for submission. Once submission is complete, the system will generate a manuscript ID and will send an email regarding your submission. Meanwhile, the authors can submit or track articles via editors@jlsb.science-line.com ; jlsb.editors@gmail.com. All manuscripts must be checked (by English native speaker) and submitted in English for evaluation (in totally confidential and impartial way).

Supplementary information

The online submission form allows supplementary information to be submitted together with the main manuscript file and covering letter. If you have more than one supplementary files, you can submit the extra ones by email after the initial [submission](#). Author guidelines are specific for each journal. Our Word template can assist you by modifying your page layout, text formatting, headings, title page, image placement, and citations/references such that they agree with the guidelines of journal. If you believe your article is fully edited per journal style, please use our [MS Word template](#) before submission. **Supplementary materials** may include figures, tables, methods, videos, and other materials. They are available online linked to the original published article. Supplementary tables and figures should be labeled with a "S", e.g. "Table S1" and "Figure S1". The maximum file size for supplementary materials is 10MB each. Please keep the files as small possible to avoid the frustrations experienced by readers with downloading large files.

Submission to the Journal is on the understanding that

- 1.The article has not been previously published in any other form and is not under consideration for publication elsewhere;
- 2.All authors have approved the submission and have obtained permission for publish work.
- 3.Researchers have proper regard for conservation and animal welfare considerations. Attention is drawn to the ['Guidelines for the Treatment of Animals in Research and Teaching'](#). Any possible adverse consequences of the work for populations or individual organisms must be weighed against the possible gains in knowledge and its practical applications. If the approval of an ethics committee is required, please provide the name of the committee and the approval number obtained.

Ethics Committee Approval

Experimental research involving human or animals should have been approved by author's institutional review board or ethics committee. This information can be mentioned in the manuscript including the name of the board/committee that gave the approval. Investigations involving humans will have been performed in accordance with the principles of [Declaration of Helsinki](#). And the use of animals in experiments will have observed the *Interdisciplinary Principles and Guidelines for the Use of Animals in Research, Testing, and Education* by the New York Academy of Sciences, Ad Hoc Animal Research Committee. If the manuscript contains photos or parts of photos of patients, informed consent from each patient should be obtained. Patient's identities and privacy should be carefully protected in the manuscript.

Graphical Abstract

Authors should provide a graphical abstract (a beautifully designed feature figure) to represent the paper aiming to catch the attention and interest of readers. Graphical abstract will be published online in the table of content. The graphical abstract should be colored, and kept within an area of 12 cm (width) x 6 cm (height) or with similar format. Image should have a minimum resolution of 300 dpi and line art 1200dpi.

Note: Height of the image should be no more than the width.

Please avoid putting too much information into the graphical abstract as it occupies only a small space. Authors can provide the graphical abstract in the format of PDF, Word, PowerPoint, jpg, or png, after a manuscript is accepted for publication. For preparing a Professional Graphical Abstract, please click [here](#).



Presentation of the article

Main Format

First page of the manuscripts must be properly identified by the title and the name(s) of the author(s). It should be typed in Times New Roman (font sizes: 17pt in capitalization for the title, 10pt for the section headings in the body of the text and the main text, double spaced, in A4 format with 2cm margins (both doc./docx formats). All pages and lines of the main text should be numbered consecutively throughout the manuscript. Abbreviations in the article title are not allowed. Manuscripts should be arranged in the following order:

- 1. TITLE** (brief, attractive and targeted)
- 2. Name(s) and Affiliation(s) of author(s)** (including post code and corresponding Email)
- 3. ABSTRACT**
- 4. Key words** (separate by semicolons; or comma,)
- 5. Abbreviations** (those used throughout the manuscript)
- 6. INTRODUCTION** (clear statement of the problem, the relevant literature on the subject, and the proposed approach or solution)
- 7. MATERIAL AND METHOD** (should be complete enough to allow experiments to be reproduced)
- 8. RESULTS**
- 9. DISCUSSION**
- 10. CONCLUSION**
- 11. DECLARATIONS** (Acknowledgements, Consent to publish, Competing interests, Authors' contributions, and Availability of data etc.)
- 12. REFERENCES**
- 13. Tables**
- 14. Figures**
- 15. Graphs**

Results and Discussion can be presented jointly.

Discussion and Conclusion can be presented jointly.

Article Sections Format

Title should be a brief phrase describing the contents of the paper. The first letter of each word in title should use upper case. The Title Page should include the author(s)'s full names and affiliations, the name of the corresponding author along with phone and e-mail information. Present address (es) of author(s) should appear as a footnote.

Abstract should be informative and completely self-explanatory, briefly present the topic, state the scope of the experiments, indicate significant data, and point out major findings and conclusions. The abstract should be 150 to 300 words in length. Complete sentences, active verbs, and the third person should be used, and the abstract should be written in the past tense. Standard nomenclature should be used and abbreviations should be avoided. No literature should be cited.

Following the abstract, about 3 to 8 **key words** that will provide indexing references should be listed.

Introduction should provide a clear statement of the problem, the relevant literature on the subject, and the proposed approach or solution. It should be understandable to colleagues from a broad range of scientific disciplines.

Material and Method should be complete enough to allow experiments to be reproduced. However, only truly new procedures should be described in detail; previously published procedures should be cited, and important modifications of published procedures should be mentioned briefly. Capitalize trade names and include the manufacturer's name and address. Subheadings should be used. Methods in general use need not be described in detail. The **ethical approval** for using human and animals in the researches should be indicated in this section with a separated title.

Results should be presented with clarity and precision. The results should be written in the past tense when describing findings in the author(s)'s experiments. Previously published findings should be written in the present tense. Results should be explained, but largely without referring to the literature. In case of the effectiveness of a particular drug or other substances as inhibitor in biological or biochemical processes, the results should be provided as **IC₅₀** (**half maximal inhibitory concentration**) or similar appropriate manner.

Discussion should interpret the findings in view of the results obtained in this and in past studies on this topic. State the conclusions in a few sentences at the end of the paper. The Results and Discussion sections can include subheadings, and when appropriate, both sections can be combined.

Conclusion should be brief and tight about the importance of the work or suggest the potential applications and extensions. This section should not be similar to the Abstract content.

Declarations including Acknowledgements, Author contribution, Competing interests, Consent to publish, and Availability of data etc.

Tables should be kept to a minimum and be designed to be as simple as possible. Tables are to be typed double-spaced throughout, including headings and footnotes. Each table should be on a separate page, numbered consecutively in Arabic numerals and supplied with a heading and a legend. Tables should be self-explanatory without reference to the text. The details of the methods used in the experiments should preferably be described in the legend instead of in the text. The same data should not be presented in both table and graph forms or repeated in the text.

Figure legends should be typed in numerical order on a separate sheet. Graphics should be prepared using applications capable of generating high resolution GIF, TIFF, JPEG or PowerPoint before pasting in the Microsoft Word manuscript file. Use Arabic numerals to designate figures and upper case letters for their parts (Figure 1). Begin each legend with a title and include sufficient description so that the figure is understandable without reading the text of the manuscript. Information given in legends should not be repeated in the text.

Declarations

Please ensure that the sections: Ethics (and consent to participate, if any), Acknowledgements, Author contribution, Competing interests, Consent to publish, Availability of data and materials are included at the end of your manuscript in a Declarations section.

Acknowledgements

We encourage authors to include an Acknowledgements section. Please acknowledge anyone who contributed towards the study by making substantial contributions to conception, design, acquisition of data, or analysis and interpretation of data, or who was involved in drafting the manuscript or revising it critically for important intellectual content, but who does not meet the criteria for authorship. Please also include their source(s) of funding. Please also acknowledge anyone who contributed materials essential for the study. Authors should obtain permission to acknowledge from all those mentioned in the Acknowledgements. Please list the source(s) of funding for the study, for each author, and for the manuscript preparation in the acknowledgements section. Authors must describe the role of the funding body, if any, in study design; in the collection, analysis, and interpretation of data; in the writing of the manuscript; and in the decision to submit the manuscript for publication.

Author contribution

For manuscripts with more than one author, JLSB require an Author Contributions section to be placed after the Acknowledgements section. An 'author' is generally considered to be someone who has made substantive intellectual contributions to a published study. To qualify as an author one should 1) have made substantial contributions to conception and design, or acquisition of data, or analysis and interpretation of data; 2) have been involved in drafting the manuscript or revising it critically for important intellectual content; and 3) have given final approval of the version to be published. Each author should have participated sufficiently in the work to take public responsibility for appropriate portions of the content. Acquisition of funding, collection of data, or general supervision of the research group, alone, does not justify authorship. **We suggest the following format/example** (please use initials to refer to each author's contribution): AB carried out the molecular genetic studies, participated in the sequence alignment and drafted the manuscript. JY carried out the immunoassays. MT participated in the sequence alignment. ES participated in the design of the study and performed the statistical analysis. FG conceived of the study, and participated in its design and coordination and helped to draft the manuscript. All authors read and approved the final manuscript.

For authors that equally participated in a study please write 'All/Both authors contributed equally to this work.' Contributors who do not meet the criteria for authorship should be listed in an acknowledgements section.

Competing interests

Competing interests that might interfere with the objective presentation of the research findings contained in the manuscript should be declared in a paragraph heading "Competing interests" (after Acknowledgment or Author Contribution sections). Examples of competing interests are ownership of stock in a company, commercial grants, board membership, etc. If there is no competing interest, please use the statement "The authors declare that they have no competing interests.".

Journal of Life Science and Biomedicine adheres to the definition of authorship set up by [The International Committee of Medical Journal Editors \(ICMJE\)](#). According to the ICMJE authorship criteria should be based on 1) substantial contributions to conception and design of, or acquisition of data or analysis and interpretation of data, 2) drafting the article or revising it critically for important intellectual content and 3) final approval of the version to be published. Authors should meet conditions 1, 2 and 3. It is a requirement that all authors have been accredited as appropriate upon submission of the manuscript. Contributors who do not qualify as authors should be mentioned under Acknowledgements.

Consent to publish

Please include a 'Consent for publication section in your manuscript. If your manuscript contains any individual person's data in any form (including individual details, images or videos), consent to publish must be obtained from that person, or in the case of children, their parent or legal guardian. All presentations of case reports must have consent to publish. You can use your institutional consent form or our consent form if you prefer. You should not send the form to us on submission, but we may request to see a copy at any stage (including after publication). If your manuscript does not contain any individual persons data, please state "Not applicable" in this section.

Change in authorship

We do not allow any change in authorship after provisional acceptance. We cannot allow any addition, deletion or change in sequence of author name. We have this policy to prevent the fraud.

Data deposition

Nucleic acid sequences, protein sequences, and atomic coordinates should be deposited in an appropriate database in time for the accession number to be included in the published article. In computational studies where the sequence information is unacceptable for inclusion in databases because of lack of experimental validation, the sequences must be published as an additional file with the article.

REFERENCES

A JLSB reference style for [EndNote](#) may be found [here](#). However, we prefer [Vancouver](#) referencing style that is often used in medicine and the natural sciences. Uniform requirements for manuscripts submitted to Biomedical Journals, published by International Committee of Medical Journal Editors, includes a list with examples of references https://www.nlm.nih.gov/bsd/uniform_requirements.html in the *Vancouver* style.

References should be numbered consecutively and cited in the text by number in square brackets [1, 2] instead of parentheses (and not by author and date). References should not be formatted as footnotes. Avoid putting personal communications and unpublished observations as references. All the cited papers in the text must be listed in References. All the papers in References must be cited in the text. Where available, URLs for the references should be provided.

Examples (at the text, blue highlighted)

Smit [1] ...; Smit and Janak [2]...; Nurai et al. [3] reported that ; ... [1], --- [2, 3], --- [3-7]. The references at the end of this document are in the preferred referencing style. Give all authors' names; do not use "et al." unless there are six authors or more. Use a space after authors' initials. Papers that have not been published should be cited as "unpublished". Papers that have been accepted for publication, but not yet specified for an issue should be cited as "to be published". Papers that have been submitted for publication should be cited as "submitted for publication". Capitalize only the first word in a paper title, except for proper nouns and element symbols. For papers published in translation journals, please give the English citation first, followed by the original foreign-language citation.

Acceptable Examples (at References section)

For Journals:

1. Hasan V, Sri Widodo M and Semedi B. 2015. Oocyte diameter distribution and fecundity of Javaen Barb (*Systemus Orphoides*) at the start of rainy season in Lenteng River, East Java, Indonesia insurance. *J. Life Sci Biomed*, 5(2): 39-42. DOI, Link
2. Karen KS, Otto CM. 2007. Pregnancy in women with valvular heart disease. *Heart*. 2007 May; 93(5): 552-558. DOI, Link
3. Doll MA, Salazar-González RA, Bodduluri S, Hein DW. Arylamine N-acetyltransferase 2 genotype-dependent N-acetylation of isoniazid in cryopreserved human hepatocytes. *Acta Pharm Sin B*, 2017; 7(4):517-522. DOI, Link

For In press manuscripts (maximum 2):

Hasan V, Sri Widodo M and Semedi B. 2015. Oocyte Diameter Distribution and Fecundity of Javaen Barb (*Systemus Orphoides*) at the Start of Rainy Season in Lenteng River, East Java, Indonesia insurance. In press.

For symposia reports and abstracts:

Cruz EM, Almatar S, Aludul EK and Al-Yaqout A. 2000. Preliminary Studies on the Performance and Feeding Behaviour of Silver Pomfret (*Pampus argentens euphrasen*) Fingerlings fed with Commercial Feed and Reared in Fibreglass Tanks. *Asian Fisheries Society Manila, Philippine* 13: 191-199. DOI, Link

For Conference:

Skinner J, Fleener B and Rinchiuso M. 2003. Examining the Relationship between Supervisors and Subordinate Feeling of Empowerment with LMX as A Possible Moderator. 24th Annual Conference for Industrial Organizational Behavior. DOI, Link

For Book:

Russell, Findlay E, 1983. Snake Venom Poisoning, 163, Great Neck, NY: Scholium International. ISBN 0-87936-015-1. DOI, Link

For Web Site:

Bhatti SA and Firkins JT. 2008. http://www.ohioline.osu.edu/sc1156_27.html. DOI, Link

Nomenclature and Abbreviations

Nomenclature should follow that given in NCBI web page and Chemical Abstracts. Standard abbreviations are preferable. If a new abbreviation is used, it should be defined at its first usage. Abbreviations should be presented in one paragraph, in the format: "term: definition". Please separate the items by ";".

E.g. ANN: artificial neural network; CFS: closed form solution; ...

Abbreviations of units should conform with those shown below:

Decilitre	dl	Kilogram	kg
Milligram	mg	hours	h
Micrometer	mm	Minutes	min
Molar	mol/L	Mililitre	ml
Percent	%	.	

Other abbreviations and symbols should follow the recommendations on units, symbols and abbreviations: in "A guide for Biological and Medical Editors and Authors (the Royal Society of Medicine London 1977). Papers that have not been published should be cited as "unpublished". Papers that have been accepted for publication, but not yet specified for an issue should be cited as "to be published". Papers that have been submitted for publication should be cited as "submitted for publication".

Formulae, numbers and symbols

1. Typewritten formulae are preferred. Subscripts and superscripts are important. Check disparities between zero (0) and the letter O, and between one (1) and the letter I.
2. Describe all symbols immediately after the equation in which they are first used.
3. For simple fractions, use the solidus (/), e.g. 10 /38.
4. Equations should be presented into parentheses on the right-hand side, in tandem.
5. Levels of statistical significance which can be used without further explanations are *P < 0.05, **P < 0.01, and ***P < 0.001.
6. In the English articles, a decimal point should be used instead of a decimal comma.
7. Use Symbol fonts for "±"; "≤" and "≥" (avoid underline).
8. In chemical formulae, valence of ions should be given, e.g. Ca²⁺ and CO₃²⁻, not as Ca⁺⁺ or CO₃.
9. Numbers up to 10 should be written in the text by words. Numbers above 1000 are recommended to be given as 10 powered x.
10. Greek letters should be explained in the margins with their names as follows: Αα - alpha, Ββ - beta, Γγ - gamma, Δδ - delta, Εε - epsilon, Ζζ - zeta, Ηη - eta, Θθ - theta, Ιι - iota, Κκ - kappa, Λλ - lambda, Μμ - mu, Νν - nu, Ξξ - xi, Οο - omicron, Ππ - pi, Ρρ - rho, Σσ - sigma, Ττ - tau, Υυ - ipsilon, Φφ - phi, Χχ - chi, Ψψ - psi, Ωω - omega. Please avoid using math equations in Word whenever possible, as they have to be replaced by images in xml full text.

Review/Decisions/Processing/Policy

Firstly, all manuscripts will be checked by [Docol@c](#), a plagiarism finding tool. The received papers with plagiarism rate of more than 30% will be rejected. Manuscripts that are judged to be of insufficient quality or unlikely to be competitive enough for publication will be returned to the authors at the initial stage. The remaining manuscripts go through a single-blind review process by external reviewers selected by section editor of JLSB, who are research workers specializing in the relevant field of study. One unfavourable review means that the paper will not be published and possible decisions are: accept as is, minor revision, major revision, or reject. The corresponding authors should submit back their revisions within 14 days in the case of minor revision, or 30 days in the case of major revision. Manuscripts with significant results are typically published at the highest priority. The editor who received the final revisions from the corresponding authors shall not be hold responsible for any mistakes shown in the final publication.

The submissions will be processed free of charge for invited authors, authors of hot papers, and corresponding authors who are editorial board members of the *Journal of Life Science and Biomedicine*. This journal encourage the academic institutions in low-income countries to publish high quality scientific results, free of charges.

Plagiarism

Manuscripts are screened for plagiarism by [Docol@c](#), before or during publication, and if found (more than 30% duplication limit) they will be rejected at any stage of processing. If we discovered accidental duplicates of published article(s) that are determined to violate our journal publishing ethics guidelines (such as multiple submission, bogus claims of authorship, plagiarism, fraudulent use of data or the like), the article will be "Withdrawn" from SCIENCELINE database. Withdrawn means that the article content (HTML and PDF) is removed and replaced with a HTML page and PDF simply stating that the article has been withdrawn according to the [Scienceline Policy](#) on Published Article Withdrawal.

Date of issue

All accepted articles are published bimonthly around 25th of January, March, May, July, September and November, each year in full text on the internet.


The OA policy

Journal of Life Science and Biomedicine is an open access journal which means that all content is freely available without charge to the user or his/her institution. Users are allowed to read, download, copy, distribute, print, search, or link to the full texts of the articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author. This is in accordance with the [BOAI definition of Open Access](#).

Submission Preparation Checklist

- Authors are required to check off their submission's compliance with all of the following items, and submissions may be returned to authors that do not adhere to the following guidelines.
- The submission has not been previously published, nor is it before another journal for consideration (or an explanation has been provided in Comments to the Editor).
- The submission file is in Microsoft Word, RTF, or PDF document file format. Where available, URLs for the references have been provided.
- The text is single-spaced; uses a 12-point font; and all illustrations, figures, and tables are placed within the text at the appropriate points, rather than at the end. The text adheres to the stylistic and bibliographic requirements outlined in the Author Guidelines.

Paper Submission Flow



SCIENCELINE PUBLISHING CORPORATION


Scienceline Publication Ltd is a limited liability non-profit non-stock corporation incorporated in Turkey, and also is registered in Iran. Scienceline journals that concurrently belong to many societies, universities and research institutes, publishes internationally peer-reviewed open access articles and believe in sharing of new scientific knowledge and vital research in the fields of life and natural sciences, animal sciences, engineering, art, linguistic, management, social and economic sciences all over the world. Scienceline journals include:

Online Journal of Animal and Feed Research



ISSN 2228-7701; Bi-monthly
[View Journal](#) | [Editorial Board](#)
 Email: editors@ojaf.r.ir
[Submit Online >>](#)

Journal of Civil Engineering and Urbanism




ISSN 2252-0430; Bi-monthly
[View Journal](#) | [Editorial Board](#)
 Email: ojceu@ojceu.ir
[Submit Online >>](#)

Journal of Life Sciences and Biomedicine




ISSN: 2251-9939; Bi-monthly
[View Journal](#) | [Editorial Board](#)
 Email: editors@jlsb.science-line.com
[Submit Online >>](#)

Asian Journal of Medical and Pharmaceutical Researches




ISSN: 2322-4789; Quarterly
[View Journal](#) | [Editorial Board](#)
 Email: editor@ajmpr.science-line.com
[Submit Online >>](#)

Journal of World's Poultry Research




ISSN: 2322-455X; Quarterly
[View Journal](#) | [Editorial Board](#)
 Email: editor@jwpr.science-line.com
[Submit Online >>](#)

World's Veterinary Journal




ISSN: 2322-4568; Quarterly
[View Journal](#) | [Editorial Board](#)
 Email: editor@wj.science-line.com
[Submit Online >>](#)

Journal of Educational and Management Studies




ISSN: 2322-4770; Quarterly
[View Journal](#) | [Editorial Board](#)
 Email: info@jems.science-line.com
[Submit Online >>](#)

Journal of World's Electrical Engineering and Technology



ISSN: 2322-5114; Irregular
[View Journal](#) | [Editorial Board](#)
 Email: editor@jweet.science-line.com
[Submit Online >>](#)

Journal of Art and Architecture Studies



ISSN: 2383-1553; Irregular
[View Journal](#) | [Editorial Board](#)
 Email: jaas@science-line.com
[Submit Online >>](#)

Asian Journal of Social and Economic Sciences



ISSN: 2383-0948; Quarterly
[View Journal](#) | [Editorial Board](#)
 Email: ajses@science-line.com
[Submit Online >>](#)

Journal of Applied Business and Finance Researches



ISSN: 2382-9907; Quarterly
[View Journal](#) | [Editorial Board](#)
 Email: jabfr@science-line.com
[Submit Online >>](#)

Scientific Journal of Mechanical and Industrial Engineering



ISSN: 2383-0980; Quarterly
[View Journal](#) | [Editorial Board](#)
 Email: sjmie@science-line.com
[Submit Online >>](#)